arXiv:hep-lat/0109022v2 3 Oct 2001

hep-lat/0109022
UTHEP-447
UTCCP-P-112

Chiral properties of domain-wall fermions from eigenvalues of 4 dimensional Wilson-Dirac operator

S. Aoki and Y. Taniguchi
Institute of Physics, University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan

(February 1, 2008)

Abstract

We investigate chiral properties of the domain-wall fermion (DWF) system by using the four-dimensional hermitian Wilson-Dirac operator. We first derive a formula which connects a chiral symmetry breaking term in the five dimensional DWF Ward-Takahashi identity with the four dimensional Wilson-Dirac operator, and simplify the formula in terms of only the eigenvalues of the operator, using an ansatz for the form of the eigenvectors. For a given distribution of the eigenvalues, we then discuss the behavior of the chiral symmetry breaking term as a function of the fifth dimensional length. We finally argue the chiral property of the DWF formulation in the limit of the infinite fifth dimensional length, in connection with spectra of the hermitian Wilson-Dirac operator in the infinite volume limit as well as in the finite volume.

Typeset using REVTEX

I. INTRODUCTION

A suitable definition of the chiral symmetry has been a long standing problem in lattice field theories. Recently an ultimate solution to this problem seems to appear in the form of the Ginsberg-Wilson relation [1,2]. Two explicit examples of the lattice fermion operators which satisfies the Ginsberg-Wilson relation have been found so far: One is the perfect lattice Dirac operator constructed via the renormalization group transformation [3,4] and the other is the overlap Dirac(OD) operator [5,6] derived from the overlap formalism [7] or from the domain-wall fermion(DWF) [8–10] in the limit of the infinite length of the 5th dimension. Since the explicit form is simpler for the latter, a lot of numerical investigations [11–18] as well as analytic considerations [19,20] have been carried out for the domain-wall fermion or the overlap Dirac fermion.

Recent numerical investigations for the domain-wall fermion [14,15], however, bring puzzling results, which are summarized as follows.

Analytic considerations suggest that the overlap or domain-wall fermion works well at sufficiently weak gauge coupling or equivalently for sufficiently smooth gauge configurations [19,20]. Initial numerical investigations supported this result [11–13]. On the other hand, further investigations indicate that the domain-wall fermion at stronger coupling ceases to describe the massless fermion even in the $N_5 \rightarrow \infty$ limit [14,15], where N_5 is the number of sites in the 5th dimension. Analytic results [21–23] in the strong coupling limit are controversial. However the latest one [24] also suggests that the DWF does not work in the limit.

It has been argued [14] that this result may be understood by the relation between the phase structure of the lattice QCD with the 4 dimensional Wilson fermion [25] and zero eigenvalues of the 4 dimensional Wilson-Dirac operator. It is well-known that the zero eigenvalues cause a trouble for the domain-wall fermion (or the overlap) [10,19,20], therefore it is natural to consider that the success/failure of DWF depends on the absence/presence of the zero eigenvalues. On the other hand, the zero eigenvalues of the 4 dimensional Wilson Dirac operator, denoted as D_W, is related to the parity-flavor breaking order parameter, $\langle \bar{q} i\gamma_5\tau^3 q\rangle$, where τ^3 is the 2×2 flavor matrix. Introducing the external source H coupled to $\bar{q} i\gamma_5\tau^3 q$, the following relation is easily derived.

$$\begin{aligned}
\left\langle \bar{q} i\gamma_5\tau^3 q\right\rangle &= -\lim_{H\rightarrow 0^+}\lim_{V\rightarrow\infty}\frac{1}{V}\mathrm{Tr}\frac{i\gamma_5\tau^3}{D_W + i\gamma_5\tau^3 H} \\
&= -\lim_{H\rightarrow 0^+}\lim_{V\rightarrow\infty}\frac{1}{V}\mathrm{tr}\left[\frac{i\gamma_5}{D_W + i\gamma_5 H} - \frac{i\gamma_5}{D_W - i\gamma_5 H}\right] \\
&= -i\lim_{H\rightarrow 0^+}\lim_{V\rightarrow\infty}\frac{1}{V}\mathrm{tr}\left[\frac{1}{H_W + iH} - \frac{1}{H_W - iH}\right] \\
&= -i\lim_{H\rightarrow 0^+}\lim_{V\rightarrow\infty}\frac{1}{V}\sum_n\left\langle \lambda_n\left|\frac{1}{\lambda_n + iH} - \frac{1}{\lambda_n - iH}\right|\lambda_n\right\rangle \\
&= -i\lim_{H\rightarrow 0^+}\int d\,\lambda\rho_{H_W}(\lambda)\left[\frac{1}{\lambda + iH} - \frac{1}{\lambda - iH}\right] \\
&= -i\int d\,\lambda\rho_{H_W}(\lambda)(-2\pi i)\delta(\lambda) = -2\pi\rho_{H_W}(0), \qquad (1.1)
\end{aligned}$$

where λ_n and $|\lambda_n\rangle$ are the eigenvalue and the eigenstate of the hermitian Wilson-Dirac operator $H_W = \gamma_5 D_W$ and $\rho_{H_W}(\lambda)$ is the density of eigenvalues of H_W, defined by

$$\rho_{H_W}(\lambda) = \lim_{V\to\infty} \frac{1}{V}\sum_n \delta(\lambda - \lambda_n). \tag{1.2}$$

The expected phase structure of lattice QCD with the Wilson fermion [25] is given in Fig. 1, where g is the gauge coupling and M is the mass parameter. In the region B, $\langle \bar{q} i\gamma_5 \tau^3 q\rangle \neq 0$, thus the density of the zero eigenvalues is nonzero. According to this phase structure, the DWF is successful between $0 < M < 2$ in the weak coupling limit of QCD. This allowed region of M agrees with the analytic consideration of DWF [9]. Once the gauge coupling becomes nonzero, the allowed region of M shrinks and moves to larger values. This property has been predicted by the mean-field analysis [26] and has numerically been observed [14]. If the coupling becomes large so that $\beta = 6/g^2 < \beta_c$, the allowed range of M disappears and the massless fermion ceases to exist in the domain-wall QCD (DWQCD).

Numerical results mentioned before [14,15] that DWF does not work in the strong coupling region seem to agree with the above expectation. This interpretation, however, has difficulties. Let us summarize the recent numerical result [14], where the quenched DWQCD has been investigated with the RG improved gauge action as well as the ordinary plaquette gauge action. With the former gauge action the quenched DWQCD works at $\beta = 2.6$ and fails at $\beta = 2.2$, suggesting that $2.2 < \beta_c < 2.6$, while with the latter gauge action the quenched DWQCD fails even at $\beta = 6.0$, indicating $\beta_c > 6.0$. One problem is that the latter condition $\beta_c > 6.0$ with the plaquette gauge contradicts the previous numerical investigation [27], which concludes that the region without parity-flavor breaking (: the allowed region in the present case) exists $\beta = 6.0$. A more serious problem is that the numerical analysis for the density of eigenvalues indicates non-zero value of $\rho_{H_W}(0)$ at any β [28] with the plaquette action, and the similar conclusion is obtained by the same analysis with the RG improved action [29]. If this is true, the phase structure in Fig.1 is incorrect in the quenched QCD: the gap of the parity-flavor breaking phase never shows up in the weak coupling region, leading to the conclusion that DWF does not work at all in the quenched QCD.

This complicated situation is summarized as follows. While being consistent with the numerical result of the DWQCD for the plaquette action [14,15], the numerical analysis for the density of the eigenvalues [28,29] contradicts the phase structure of QCD with Wilson fermion [25,27] and the numerical result of DWQCD with the RG action [14]. Furthermore the numerical result of the DWQCD with the plaquette action seems apparently inconsistent with the numerical result for the phase structure of the Wilson fermion with the same gauge action.

In this paper, we try to resolve the above mutual inconsistency, by analysing eigenvalues of the 4 dimensional hermitian Wilson-Dirac operator. In Sec. II the action and the axial Ward-Takahashi identity are given for the latter use. In Sec. III, we derive the formula for the explicit chiral symmetry breaking term of DWQCD, m_{5q}, in terms of eigenvalues and eigenstates of the modified hermitian Wilson-Dirac operator. In Sec. IV, assuming the form of the eigenstates, the formula derived in Sec. III is reduced to a much simpler expression in terms of the eigenvalues only. The dependence of m_{5q} on N_5 is discussed in Sec. V, for a given model of the density of the eigenvalues. In Sec. VI, using the simplified formula, we discuss the chiral properties of DWQCD first in the finite volume, and consider it in the infinite volume limit. Some comments are also given on the phase structure and the exceptional configurations in the lattice QCD with the Wilson fermion. Our conclusion is presented in Sec. VII.

II. ACTION AND AXIAL WARD-TAKAHASHI IDENTITY

We employ Shamir's domain-wall fermion action [9,10]. Flipping the sign of the Wilson term and the domain wall height M, we write

$$S_{DW} = a^4 \sum_{x,y,s,s'} \overline{\psi}(x,s)\left[\left(D_W - \frac{M}{a}\right)_{x,y} \delta_{s,s'} + (D_5)_{s,s'}\,\delta_{x,y}\right]\psi(y,s'), \tag{2.1}$$

$$D_W = \sum_{\mu}\left(\gamma_\mu \frac{1}{2}\left(\nabla_\mu^f + \nabla_\mu^b\right) - \frac{a}{2}\nabla_\mu^f \nabla_\mu^b\right), \tag{2.2}$$

$$D_5 = \gamma_5 \frac{1}{2}\left(\nabla_5^f + \nabla_5^b\right) - \frac{a_5}{2}\nabla_5^f \nabla_5^b, \tag{2.3}$$

where x, y are four-dimensional space-time coordinates, and s, s' are fifth-dimensional or "flavor" indices, bounded as $1 \le s, s' \le N_5$ with the free boundary condition at both ends (we assume N_5 to be even). The domain-wall height M should be taken as $0 < M < 2$ at tree level to give massless fermion modes. ∇_μ^f and ∇_μ^b are forward and backward derivative in four dimensions

$$\left(\nabla_\mu^f\right)_{x,y} = \frac{1}{a}\left(\delta_{x+\mu,y} U_\mu(x) - \delta_{x,y}\right), \tag{2.4}$$

$$\left(\nabla_\mu^b\right)_{x,y} = \frac{1}{a}\left(\delta_{x,y} - \delta_{x-\mu,y} U_\mu^\dagger(y)\right) \tag{2.5}$$

and ∇_5^f and ∇_5^b are those in the fifth dimension with the boundary condition

$$\left(\nabla_5^f\right)_{s,s'} = \frac{1}{a_5}\begin{cases} \delta_{s+1,s'} - \delta_{s,s'} & (1 \le s < N_5) \\ a_5 m_f \delta_{s',1} - \delta_{s,s'} & (s = N_5) \end{cases}, \tag{2.6}$$

$$\left(\nabla_5^b\right)_{s,s'} = \frac{1}{a_5}\begin{cases} \delta_{s,s'} - a_5 m_f \delta_{s',N_5} & (s = 1) \\ \delta_{s,s'} - \delta_{s-1,s'} & (1 < s \le N_5) \end{cases}, \tag{2.7}$$

where m_f is the mass for the quark field. The light fermion mode is extracted by the 4-dimensional quark field defined on two boundaries of the fifth dimension,

$$\begin{aligned} q(x) &= P_L \psi(x,1) + P_R \psi(x,N_5), \\ \overline{q}(x) &= \overline{\psi}(x,N_5) P_L + \overline{\psi}(x,1) P_R, \end{aligned} \tag{2.8}$$

where $P_{R/L}$ is the projection matrix $P_{R/L} = (1 \pm \gamma_5)/2$. The quark mass term induces a coupling between these boundary fields through the bare quark mass m_f.

In this paper we investigate chiral property of the domain-wall fermion through the axial Ward-Takahashi (WT) identity defined in Ref. [10] for the non-singlet axial transformation

$$\delta_A^a \psi(x,s) = i\epsilon(N_5 + 1 - 2s) T^a \psi(x,s), \tag{2.9}$$

$$\delta_A^a \overline{\psi}(x,s) = -i\epsilon(N_5 + 1 - 2s) T^a \overline{\psi}(x,s), \tag{2.10}$$

where $\epsilon(x)$ is a sign function of x. The WT identity for some operator $\mathcal{O}$ is written as follows.

$$\left\langle \nabla_\mu^b A_\mu^a(x)\mathcal{O}\right\rangle = 2m_f \left\langle P^a(x)\mathcal{O}\right\rangle + 2\left\langle J_{5q}^a(x)\mathcal{O}\right\rangle - \left\langle \delta_A^a \mathcal{O}\right\rangle , \tag{2.11}$$

where an axial vector current A_μ^a, a pseudo scalar density P^a, and an explicit chiral symmetry breaking term J_{5q} at finite N_5 are given by

$$\begin{aligned} A_\mu^a(x) &= \sum_{s=1}^{N_5} \epsilon(N_5+1-2s)\frac{1}{2}\Big(\overline{\psi}(x,s)T^a(1-\gamma_\mu)U_\mu(x)\psi(x+\mu,s) \\ &\qquad -\overline{\psi}(x+\mu,s)(1+\gamma_\mu)U_\mu^\dagger(x)T^a\psi(x,s)\Big), & (2.12) \\ P^a(x) &= \overline{q}(x)\gamma_5 T^a q(x), & (2.13) \\ J_{5q}^a(x) &= \frac{1}{a_5}\left(\overline{\psi}(x,\frac{N_5}{2}+1)T^a P_R \psi(x,\frac{N_5}{2}) - \overline{\psi}(x,\frac{N_5}{2})T^a P_L \psi(x,\frac{N_5}{2}+1)\right) \\ &= \frac{1}{a_5}\overline{\psi}'(x,\frac{N_5}{2})T^a\gamma_5\psi'(x,\frac{N_5}{2}). & (2.14) \end{aligned}$$

Here we define the following fields according to [6]

$$\begin{aligned} \psi'(x,\frac{N_5}{2}) &= P_R\psi(x,\frac{N_5}{2}) + P_L\psi(x,\frac{N_5}{2}+1), & (2.15) \\ \overline{\psi}'(x,\frac{N_5}{2}) &= \overline{\psi}(x,\frac{N_5}{2}+1)P_R + \overline{\psi}(x,\frac{N_5}{2})P_L. & (2.16) \end{aligned}$$

In this paper we consider the identity with $\mathcal{O} = P^b(y)$

$$\begin{aligned} \left\langle \nabla_\mu^b A_\mu^a(x)P^b(y)\right\rangle &= 2m_f\left\langle P^a(x)P^b(y)\right\rangle + 2\left\langle J_{5q}^a(x)P^b(y)\right\rangle \\ &\quad -\frac{1}{a^4}\delta_{x,y}\left\langle \overline{q}(y)\{T^a,T^b\}q(y)\right\rangle \end{aligned} \tag{2.17}$$

and measure the chiral symmetry breaking effect by

$$m_{5q} = \lim_{t\to\infty}\frac{\sum_{\vec{x}}\left\langle J_{5q}(t,\vec{x})P(0)\right\rangle}{\sum_{\vec{x}}\left\langle P(t,\vec{x})P(0)\right\rangle}, \tag{2.18}$$

which we call an 'anomalous quark mass' [14]. Please notice that we have omitted flavor indices since the flavor factors are canceled in the definition of m_{5q}.

III. EFFECTIVE THEORY OF DOMAIN-WALL FERMIONS

We now derive the effective theory of the DWF system by integrating out the heavy bulk modes according to Ref. [6]. Our aim is to rewrite the numerator and denominator of the anomalous quark mass (2.18) in terms of four dimensional quantities and hence to relate m_{5q} to an hermitian Wilson-Dirac operator in the four dimensional theory. Introducing source fields to $q(x)$, $\overline{q}(x)$, $\psi'(x,N_5/2)$, $\overline{\psi}'(x,N_5/2)$, the propagators necessary for our purpose are derived as [6]

$$\langle q(x)\overline{q}(y)\rangle = \frac{a_5}{a^5}\frac{1}{1-am_f}\left(D_{N_5}^{-1}-a\right), \tag{3.1}$$

$$\left\langle \psi'(x,\frac{N_5}{2})\overline{q}(y)\right\rangle = \frac{a_5}{a^5}\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}}D_{N_5}^{-1}, \tag{3.2}$$

$$\left\langle q(x)\overline{\psi}'(y,\frac{N_5}{2})\right\rangle = \frac{a_5}{a^5}D_{N_5}^{-1}\gamma_5\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}}\gamma_5, \tag{3.3}$$

where $\widetilde{H}$ is a DWF Hamiltonian in the 5th direction, with which the transfer matrix in fifth direction is given by

$$T = e^{-a_5\widetilde{H}}. \tag{3.4}$$

D_{N_5} is a truncated Ginsparg-Wilson (GW) Dirac operator which satisfies GW relation in $N_5 \to \infty$ limit. $\widetilde{H}$ and D_{N_5} are related to the four dimensional hermitian Wilson-Dirac operator H_W as,

$$D_{N_5} = \frac{1}{2a}\left[(1+am_f)+(1-am_f)\gamma_5\tanh\frac{N_5}{2}a_5\widetilde{H}\right], \tag{3.5}$$

$$\widetilde{H} = \frac{1}{a_5}\log\frac{1+aH'}{1-aH'}, \tag{3.6}$$

$$H' = H_W\frac{1}{2+a\gamma_5H_W}, \tag{3.7}$$

$$H_W = \gamma_5\left(D_W-\frac{M}{a}\right). \tag{3.8}$$

Here we adopted Boriçi's notation [30] for $\widetilde{H}$.

The numerator of m_{5q} can be written in terms of the above propagators,

$$\begin{aligned} X(t) &= \sum_{\vec{x}}\langle J_{5q}(\vec{x},t)P(\vec{y},0)\rangle \\ &= -\frac{1}{a_5}\sum_{\vec{x}}\mathrm{tr}\left[\gamma_5\left\langle\psi'(\vec{x},t,\frac{N_5}{2})\overline{q}(\vec{y},0)\right\rangle\gamma_5\left\langle q(\vec{y},0)\overline{\psi}'(\vec{x},t,\frac{N_5}{2})\right\rangle\right] \\ &= -\frac{a_5}{a^{10}}\sum_{\vec{x},\alpha,a}\sum_{\beta,b}\left\langle I\left|\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}}D_{N_5}^{-1}\right|J\right\rangle\left\langle J\left|\left(D_{N_5}^{-1}\right)^\dagger\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}}\right|I\right\rangle, \end{aligned} \tag{3.9}$$

where $I=(\vec{x},t,\alpha,a)$, $J=(\vec{y},0,\beta,b)$. α,β are spinor indices and a,b are color indices. $|I\rangle$ is an eigen-ket in the coordinate, spinor and color spaces. In the last equality we use a relation that $\gamma_5(D_{N_5}^{-1})\gamma_5 = (D_{N_5}^{-1})^\dagger$. The denominator of m_{5q} is given by

$$\begin{aligned} Y(t) &= \sum_{\vec{x}}\langle P(\vec{x},t)P(\vec{y},0)\rangle \\ &= -\sum_{\vec{x}}\mathrm{tr}\left[\gamma_5\langle q(\vec{x},t)\overline{q}(\vec{y},0)\rangle\gamma_5\langle q(\vec{y},0)\overline{q}(\vec{x},t)\rangle\right] \\ &= -\frac{a_5^2}{a^{10}}\frac{1}{(1-am_f)^2}\sum_{\vec{x},\alpha,a}\sum_{\beta,b}\left\langle I\left|\left(D_{N_5}^{-1}-a\right)\right|J\right\rangle\left\langle J\left|\gamma_5\left(D_{N_5}^{-1}-a\right)\gamma_5\right|I\right\rangle \\ &= -\frac{a_5^2}{a^{10}}\frac{1}{(1-am_f)^2}\sum_{\vec{x},\alpha,a}\sum_{\beta,b}\left\langle I\left|D_{N_5}^{-1}\right|J\right\rangle\left\langle J\left|\left(D_{N_5}^{-1}\right)^\dagger\right|I\right\rangle. \end{aligned} \tag{3.10}$$

Note that $\langle I|J\rangle = 0$ for $t \neq 0$.

Hereafter we set the bare quark mass $m_f = 0$ without loss of generality.

IV. EXPANSION IN TERMS OF EIGENSTATES

We expand $X(t)$ and $Y(t)$ in terms of eigenstates of the hermitian operator H'. Since the numerator (3.9) has a suppression factor $1/(2\cosh\frac{N_5}{2}a_5\widetilde{H})$ only small eigenvalues of $\widetilde{H}$, or equivalently those of H', contribute to m_{5q} at large N_5. In this case we can expand H' perturbatively in terms of the four dimensional hermitian Wilson-Dirac operator H_W and can derive the formula which connect m_{5q} and the eigenvalue of H_W at large N_5.

The expansion of $X(t)$ and $Y(t)$ with these eigenstates is done by inserting a complete set of eigenstates

$$1 = \sum_n |n\rangle\langle n|\,, \tag{4.1}$$

where $|n\rangle$ is n-th eigenstate of H'

$$H'|n\rangle = \lambda'_n|n\rangle\,. \tag{4.2}$$

With this substitution $X(t)$ and $Y(t)$ are written in terms of eigenvalue and eigenfunction.

In order to further simplify these Green functions as a function of eigenvalue only, we employ the following assumptions and approximations for typical form of the eigenfunctions. The recent numerical analyses indicate that the eigenvalues of H_W seem to be classified into two groups [19,31]; one is a group of small isolated eigenvalues and the other is a group of almost continuous eigenvalues above the isolated ones. The eigenvectors associated with the isolated eigenvalues are exponentially localized at some center in the coordinate space [19,31], and those for the continuous eigenvalues are rapidly oscillating plane-wave functions in the coordinate space [31]. This property of eigenvalues is expected to hold also for H', since H' can be expanded in terms of H_W perturbatively for small eigenvalues. ¿From this consideration we assume that the eigenvector space of H' is divided into two subspaces $\mathcal{S}$ and $\widetilde{\mathcal{S}}$, where $\mathcal{S}$ is spanned by localized eigenvectors and $\widetilde{\mathcal{S}}$ is expanded with plane wave functions.

A. Localized eigenvectors

We adopt two types of the approximation for typical form of the localized eigenvectors; completely localized ones and partially localized ones. The completely localized eigenvector means that the eigenfunction ψ_n of the n-th eigenvalue has non-zero value at a single parameter (x_n, α_n, a_n)

$$\psi_n(x,\alpha,a) = \delta_{x,x_n}\delta_{\alpha,\alpha_n}\delta_{a,a_n}, \tag{4.3}$$

which gives a unit vector in (x,α,a) space. $\psi_n(x,\alpha,a)$ is given by using the n-th eigenstate $|n\rangle$ of an eigenvalue λ'_n

$$\psi_n(x,\alpha,a) = \langle x,\alpha,a|n\rangle\,, \tag{4.4}$$

$$H'\,|n\rangle = \lambda'_n\,|n\rangle\,, \tag{4.5}$$

$$\widetilde{H}\,|n\rangle = \widetilde{\lambda}(\lambda'_n)\,|n\rangle\,, \tag{4.6}$$

$$\widetilde{\lambda}(\lambda'_n) = \frac{1}{a_5}\log\frac{1+a\lambda'_n}{1-a\lambda'_n}. \tag{4.7}$$

We assume that eigenvectors with different eigenvalues reside at different points

$$\langle n|x,\alpha,a\rangle\,\langle x,\alpha,a|m\rangle = 0 \quad \text{for } n \neq m. \tag{4.8}$$

In this case $\mathcal{S}$ is spanned by a set of basis vectors

$$\begin{pmatrix}1\\0\\ \vdots\end{pmatrix}, \quad \begin{pmatrix}0\\1\\ \vdots\end{pmatrix}, \quad \cdots. \tag{4.9}$$

On the other hand, the partially localized eigenvector is non-zero at small range of (x,α,a) space. We call this small range of volume h_n as $\mathcal{S}_n$ and $\mathcal{S} = \cup_n \mathcal{S}_n$. $\mathcal{S}_n$ is spanned by a set of eigenvectors $\psi_n^{(i)}$, where i runs as $i = 1,\cdots,h_n$ and $\psi_n^{(i)}$ is given by

$$\psi_n^{(i)}(x,\alpha,a) = \langle x,\alpha,a|n,i\rangle\,, \tag{4.10}$$

$$H'\,|n,i\rangle = \lambda_n^{\prime(i)}\,|n,i\rangle\,. \tag{4.11}$$

We assume that $\mathcal{S}_n$ with different n has no overlap each other;

$$\langle m,i|x,\alpha,a\rangle\,\langle x,\alpha,a|n,j\rangle = 0 \tag{4.12}$$

for $m \neq n$ for any i,j. In the limit of $h_n \to 1$ the partially localized eigenstate is reduced to the completely localized one.

Correspondence between the eigenvalues and eigenvectors of H' and H_W is given perturbatively for small eigenvalues. We expand H' in terms of H_W

$$2H' = H_W + H_W\sum_{k=1}^{\infty}(-1)^k\left(\frac{a\gamma_5 H_W}{2}\right)^k, \tag{4.13}$$

and the eigenvalue λ_n^W and eigenstate $|n_W\rangle$ of H_W is defined by

$$H_W\,|n_W\rangle = \lambda_n^W\,|n_W\rangle\,. \tag{4.14}$$

The standard perturbation theory gives the relation between λ_n, $|n\rangle$ and λ_n^W, $|n_W\rangle$ as follows.

$$2\lambda'_n = \lambda_n^W - \frac{1}{2}\left(\lambda_n^W\right)^2\langle n_W|\gamma_5|\,n_W\rangle + \mathcal{O}(\lambda^3), \tag{4.15}$$

$$|n\rangle = |n_W\rangle - \frac{1}{2}\lambda_n^W\frac{\phi_n}{\lambda_n^W - H_W}H_W\gamma_5\,|n_W\rangle + \mathcal{O}(\lambda^3), \tag{4.16}$$

where ϕ_n is a projection operator onto the space perpendicular to the n-th eigenstate:

$$\phi_n = 1 - |n_W\rangle\,\langle n_W|\,. \tag{4.17}$$

B. Plane-wave eigenvector

We approximate the plane-wave function with that of the free theory but relax the condition for possible momenta p in the finite box. The operator H' of the free theory in momentum space is given by

$$H'(p) = \frac{1}{a}\gamma_5 \left(i\sum_\mu \gamma_\mu A_\mu(p) + B(p) \right), \tag{4.18}$$

$$A_\mu(p) = \frac{2s_\mu}{s^2 + (2 + 2\hat{s}^2 - M)^2}, \tag{4.19}$$

$$B(p) = \frac{s^2 + (2\hat{s}^2 - M)(2 + 2\hat{s}^2 - M)}{s^2 + (2 + 2\hat{s}^2 - M)^2}, \tag{4.20}$$

where $s_\mu = \sin(ap_\mu)$, $\hat{s}^2 = \sum_\mu \sin^2(ap_\mu/2)$ and $s^2 = \sum_\mu s_\mu^2$. The eigenstate of free H' and $\widetilde{H}$ becomes

$$H' \left| p, s, a \right\rangle = \lambda'(p) \left| p, s, a \right\rangle, \tag{4.21}$$

$$\widetilde{H} \left| p, s, a \right\rangle = \widetilde{\lambda}(p) \left| p, s, a \right\rangle, \tag{4.22}$$

$$\lambda'(p) = \pm\frac{1}{a}\sqrt{A^2(p) + B^2(p)}, \tag{4.23}$$

$$\widetilde{\lambda}(p) = \frac{1}{a_5} \log \frac{1 + a\lambda'(p)}{1 - a\lambda'(p)}, \tag{4.24}$$

where the spinor index s runs as $s = 1, 2, 3, 4$, and the eigenstate is degenerate in the color index a. The eigenvector in (x, α, a) space can be written as

$$\langle x, \alpha, a | p, s, b \rangle = \frac{1}{\sqrt{N_p}} U_\alpha(p, s) e^{ipx} \delta_{a,b} \tag{4.25}$$

within the subspace $\widetilde{\mathcal{S}}$, where $U_\alpha(p, s)$ is a normalized eigenfunction of free $H'(p)$, given in Appendix A, and N_p is the total number of the plane-wave eigenvectors. The free truncated overlap Dirac operator and its inverse in momentum space are given by

$$\langle k, s, a \left| D_{N_5} \right| p, t, b \rangle = \frac{1}{2a} \left(i(\gamma_\mu)_{s,t} C_\mu(p) + \delta_{s,t} E(p) \right) \delta_{k,p} \delta_{a,b}, \tag{4.26}$$

$$\left\langle k, s, a \left| D_{N_5}^{-1} \right| p, t, b \right\rangle = 2a \frac{-i(\gamma_\mu)_{s,t} C_\mu(p) + \delta_{s,t} E(p)}{C^2 + E^2} \delta_{k,p} \delta_{a,b}, \tag{4.27}$$

where

$$C_\mu(p) = \frac{A_\mu(p)}{a|\lambda'(p)|} \tanh \frac{N_5}{2} a_5 |\widetilde{\lambda}(p)|, \tag{4.28}$$

$$E(p) = \left(1 + \frac{B(p)}{a|\lambda'(p)|} \tanh \frac{N_5}{2} a_5 |\widetilde{\lambda}(p)| \right). \tag{4.29}$$

In addition we also assume that the overlap of two eigenfunctions from different groups vanishes:

$$\langle n|x,\alpha,a\rangle \langle x,\alpha,a|p,s,b\rangle = 0. \tag{4.30}$$

The eigenvalues of the hermitian Wilson-Dirac operator H_W is also given as a function of p_μ for the free theory. The free H_W in the momentum space is given by

$$H_W = \gamma_5 \left(i\gamma_\mu s_\mu + 2\hat{s}^2 - M \right). \tag{4.31}$$

The eigenvalue $\lambda_W(p)$ of free H_W is given by

$$H_W \left|\lambda_W(p)\right\rangle = \lambda_W(p) \left|\lambda_W(p)\right\rangle, \tag{4.32}$$

$$\lambda_W(p) = \pm\sqrt{s^2 + (2\hat{s}^2 - M)^2}, \tag{4.33}$$

where $|\lambda_W(p)\rangle$ is the corresponding eigenstate. Although the eigenstates of $\lambda'(p)$ and $\lambda_W(p)$ are different, $\lambda'(p)$ of H' is given in terms of $\lambda_W(p)$ and the momentum p as

$$\lambda'(p) = \frac{\lambda_W(p)}{\sqrt{\lambda_W(p)^2 + 4(2\hat{s}^2 - M) + 4}}. \tag{4.34}$$

C. Simplification of the formula of m_{5q}

We first consider a system with N_l completely localized eigenvectors and plane wave functions with N_p degrees of freedom for the momentum p. N_l and N_p satisfy a relation $N_l + 4N_cN_p = 4N_cN_x$, where N_x is a number of total sites of our lattice $N_x = n_xn_yn_zn_t$. Since the total eigenvector space is separated into two subspaces $\mathcal{S}$ and $\widetilde{\mathcal{S}}$, the complete set of this system is given by

$$1 = \sum_{n=1}^{N_l} |n\rangle \langle n| + \sum_{p=1}^{N_p} \sum_{s=1}^{4} \sum_{a=1}^{N_c} |p,s,a\rangle \langle p,s,a|, \tag{4.35}$$

where $|n\rangle \in \mathcal{S}$ and $|p,s,a\rangle \in \widetilde{\mathcal{S}}$.

We expand the numerator of m_{5q} with the complete set in the above.

$$\begin{aligned} X(t) &= -\frac{a_5}{a^{10}} \sum_{\vec{x},\alpha,a} \sum_{b,\beta} \left\langle I \left| \frac{1}{2\cosh \frac{N_5}{2} a_5 \widetilde{H}} D_{N_5}^{-1} \right| J \right\rangle \left\langle J \left| \left(D_{N_5}^{-1}\right)^\dagger \frac{1}{2\cosh \frac{N_5}{2} a_5 \widetilde{H}} \right| I \right\rangle \\ &= X_l(t) + X_c(t), \end{aligned} \tag{4.36}$$

where $X_l(t)$ and $X_c(t)$ are the contributions from the localized eigenvectors and the plane wave function, respectively. After a little algebra, the detail of which is given in Appendix B, we obtain

$$X_l(t) = \frac{1}{a_5} Y(t) \frac{1}{4N_cN_x} \sum_{n=1}^{N_l} \left(\frac{1}{2\cosh \frac{N_5}{2} a_5 \widetilde{\lambda}(\lambda'_n)} \right)^2 \tag{4.37}$$

for the contribution from the localized eigenvectors, where $Y(t)$ is the denominator of m_{5q}.

Similarly, taking $t \to \infty$ limit, we have for the contribution of the continuous eigenvalues

$$X_c(t)|_{t\to\infty} = -\frac{a_5}{a^8} n'_{xyz} \frac{1}{N_p} \sum_p \left(\frac{1}{2\cosh \frac{N_5}{2} a_5 \widetilde{\lambda}(p)} \right)^2 F(p), \tag{4.38}$$

where n'_{xyz} is a volume of $\widetilde{\mathcal{S}}$ in xyz space, and $F(p)$ is defined as

$$F(p) = \frac{4N_c}{N_p} \left(\frac{4}{C^2(p) + E^2(p)} \right). \tag{4.39}$$

We next evaluate the denominator of m_{5q} in the similar manner:

$$Y(t) = -\frac{a_5^2}{a^{10}} \sum_{\vec{x},\alpha,a} \sum_{\beta,b} \left\langle I \left| D_{N_5}^{-1} \right| J \right\rangle \left\langle J \left| \left(D_{N_5}^{-1} \right)^\dagger \right| I \right\rangle = Y_l(t) + Y_c(t), \tag{4.40}$$

where

$$Y_l(t) = \frac{N_l}{4N_c N_x} Y(t) \tag{4.41}$$

and

$$Y_c(t)|_{t\to\infty} = -\frac{a_5^2}{a^{10}} n'_{xyz} \frac{1}{N_p} \sum_p F(p). \tag{4.42}$$

These results imply

$$Y(t \to \infty) = -\frac{4N_c N_x}{4N_c N_x - N_l} \frac{a_5^2}{a^{10}} n'_{xyz} \frac{1}{N_p} \sum_p F(p). \tag{4.43}$$

We can directly see that the function $F(p)$ is almost constant for all the range of p as in Fig. 2 except some discrete points of p. Therefore it is a reasonable approximation to assume that $F(p)$ is independent of p. Combining these formula and approximations the anomalous quark mass becomes

$$\begin{aligned} m_{5q} &= \frac{\lim_{t\to\infty} X(t)}{\lim_{t\to\infty} Y(t)} \\ &= \frac{1}{a_5} \frac{1}{4N_c N_x} \left(\sum_n \left(\frac{1}{2\cosh \frac{N_5}{2} a_5 \widetilde{\lambda}(\lambda'_n)} \right)^2 + \sum_n \rho_n \left(\frac{1}{2\cosh \frac{N_5}{2} a_5 \widetilde{\lambda}_n} \right)^2 \right), \end{aligned} \tag{4.44}$$

where ρ_n is a number of degeneracy in $\widetilde{\lambda}_n$,

$$\rho_n = \sum_{p,\alpha,a} \delta_{\widetilde{\lambda}_n, \widetilde{\lambda}(p)}. \tag{4.45}$$

If the degeneracy of ρ_n in the free case is resolved by the presence of gauge fields, we are able to reconstruct m_{5q} by simply summing up all eigenvalues of H',

$$m_{5q} = \frac{1}{a_5}\frac{1}{4N_cN_x}\sum_n \left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda'_n)}\right)^2. \tag{4.46}$$

The above result can be generalized to the case with the partially localized eigenstates. The formula becomes

$$\begin{aligned} m_{5q} &= \frac{\lim_{t\to\infty} X(t)}{\lim_{t\to\infty} Y(t)} \\ &= \frac{1}{a_5}\frac{1}{4N_cN_x}\left(\sum_{n=1}^{N_l}\sum_{i=1}^{h_n}\tilde{h}_n^{(i)}\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda_n^{'(i)})}\right)^2 + \sum_n \rho_n\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}_n}\right)^2\right), \end{aligned} \tag{4.47}$$

where the n-th set of localized eigenvectors have the (local) support with the dimensionless volume of h_n, and the $\tilde{h}_n^{(i)}$ $(1 \le \tilde{h}_n^{(i)} \le 4^4 h_n)$ is the enhancement factor for the set, which depends on the shape of eigenvectors. See the detail of the derivation in Appendix C. When the degeneracy in the continuous eigenvalues is resolved with gauge fields we have

$$m_{5q}a_5 = \frac{1}{4N_cN_x}\left(\sum_{\text{local}}\tilde{h}_n\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda'_n)}\right)^2 + \sum_{\text{continuous}}\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda'_n)}\right)^2\right). \tag{4.48}$$

V. MODEL FOR EIGENVALUE DENSITY

In this section we consider the relation between the asymptotic behavior of m_{5q} in N_5 and the distribution of the continuous eigenvalues. For simplicity we neglect the effect of the localized eigenvalues here and will discuss their effect in the next section. In this simple situation we can write m_{5q} as an integral in continuous eigenvalues

$$m_{5q}a_5 = \int_{\lambda_{\min}}^{\lambda_{\max}} d\lambda\rho(\lambda)\left(\frac{1}{2\cosh\frac{N_5}{2}\lambda}\right)^2, \tag{5.1}$$

where λ is a dimensionless eigenvalue $\lambda = a_5\widetilde{\lambda}$, $\lambda_{\min}$ and $\lambda_{\max}$ are minimum and maximum of eigenvalues $0 \le \lambda_{\min} \le \lambda \le \lambda_{\max}$. Without loss of generality, we consider non-negative λ only, by taking $\rho(\lambda) = \rho_n(\lambda) + \rho_n(-\lambda)$, since $\cosh\frac{N_5}{2}\lambda$ is the even function. We adopt the following three types of $\rho(\lambda)$

$$\rho(\lambda) = \sqrt{R^2 - (\lambda - R - \delta)^2} \tag{5.2}$$

with

1. $\delta < 0$, $\lambda_{\min} = 0$, $\lambda_{\max} = 2R - |\delta|$, $\rho(0) \neq 0$,
2. $\delta = 0$, $\lambda_{\min} = 0$, $\lambda_{\max} = 2R$, $\rho(0) = 0$,

3. $\delta > 0$, $\lambda_{\min} = \delta$, $\lambda_{\max} = 2R + \delta$, $\rho(0) = 0$.

A typical form of $\rho(\lambda)$ for each case is given in Fig. 3

A main support of the integral at large N_5 resides near $\lambda_{\min}$ and asymptotic behavior of m_{5q} is evaluated by expanding $\rho(\lambda)$ around $\lambda_{\min}$ and adopting the leading term. Since contribution from the larger eigenvalues is negligible we set the integral range to $[\lambda_{\min}, \infty]$. The integral in λ is easily calculated with the following two formulas,

$$\left(\frac{1}{2\cosh\frac{N_5}{2}\lambda}\right)^2 = \sum_{n=1}^{\infty}(-)^{n-1} n e^{-nN_5\lambda}, \tag{5.3}$$

$$\int_0^{\infty} d\lambda e^{-x\lambda}\lambda^{\alpha} = \frac{\Gamma(\alpha+1)}{x^{\alpha+1}}, \tag{5.4}$$

which is valid for $x > 0$, $\alpha > -1$.

For $\delta < 0$ the asymptotic behavior of the anomalous quark mass becomes

$$m_{5q} \longrightarrow \frac{\rho(0)}{2N_5} + \mathcal{O}\left(\frac{1}{N_5^2}\right). \tag{5.5}$$

For $\delta = 0$

$$m_{5q} \longrightarrow \sqrt{\frac{\pi R}{2}}(1-\sqrt{2})\zeta\left(\frac{1}{2}\right)\frac{1}{N_5^{3/2}} + \mathcal{O}\left(\frac{1}{N_5^{5/2}}\right), \tag{5.6}$$

where ζ is the Riemann's zeta function. For $\delta > 0$ we have

$$m_{5q} \longrightarrow e^{-\delta N_5}\left(\sqrt{\frac{\pi R}{2}}\frac{1}{N_5^{3/2}} + \mathcal{O}\left(\frac{1}{N_5^{5/2}}\right)\right). \tag{5.7}$$

We can see that $\delta = 0$ gives power law even if the zero mode density vanishes $\rho(0) = 0$. We need a gap at $\lambda = 0$ in the continuous eigenvalues to realize exponential decay. Note that the similar analysis is also made in Ref. [32]. Typical form of m_{5q} is given in Fig. 4 for $R = 4$ and $\delta = -0.5$, $\delta = 0$, $\delta = 0.5$.

VI. EFFECT OF LOCALIZED EIGENSTATES AND THE INFINITE VOLUME LIMIT OF THE SYSTEM

In this section, using the formula for m_{5q} in terms of the eigenvalues, we propose our interpretation, which resolves the inconsistency among the numerical simulations for the domain-wall QCD, mentioned in the introduction.

We consider the situation that the density of the continuous eigenvalues has a gap at zero: $\rho(\lambda) = 0$ for $|\lambda| \leq \delta$ with $\delta > 0$. On the other hand, we do not put such a restriction on the localized eigenvalues, so that they can become almost zero. In this situation, the contribution of the continuous eigenvalues to m_{5q} vanishes exponentially in N_5. In the following subsections we will consider the contribution from the localized eigenstates to m_{5q}.

A. $N_5 \to \infty$ at finite volume

In this subsection we discuss the behavior of m_{5q} in the large N_5 at finite volume. This situation often corresponds to the one encountered in the numerical simulations, and in the infinite N_5 limit the domain-wall fermion becomes the overlap Dirac operator, which satisfies the Ginsparg-Wilson relation.

In the finite volume, it is almost impossible to have an *exact* zero eigenvalue. More precisely a probability to have the exact zero eigenvalue is zero, since no symmetry assures the existence of it. Indeed no exact zero is numerically found in the evaluations of eigenvalues of H_W. On the other hand, *almost* zero eigenvalues, $|\lambda| \simeq 10^{-2}$ or less, appear, and they become smaller as the volume increases. Therefore it may be reasonable to assume that the average of the smallest eigenvalue vanishes in some power of $1/V$: $\langle|\lambda_{\rm min}|\rangle = c_0 V^{-c_1}$ with $c_0, c_1 > 0$.

In this situation m_{5q} vanishes exponentially as N_5 increase:

$$m_{5q} \propto \frac{1}{4N_c V}\exp[-\frac{N_5 c_0}{V^{c_1}}] \tag{6.1}$$

as $N_5 \to \infty$. This means that DWQCD always works in the finite volume: m_{5q} vanishes exponentially in N_5. In other words, as long as the continuous eigenvalues have a gap around zero, DWF in the $N_5 \to \infty$ limit, or equivalently, the overlap Dirac fermion, works well to describe the chiral modes in the finite volume, where the smallest eigenvalue of the localized modes is small but non-zero.

According to this consideration, we speculate how DWQCD behaves as the coupling constant varies. In the strong coupling, even the continuous eigenvalues have no gap such that $|\lambda_{\rm min}| = 0$, and therefore m_{5q} vanishes only in some power of $1/N_5$. DWQCD does not work in the strong coupling region. Once the gap in the continuous eigenvalues opens ($|\lambda_{\rm min}| > \delta > 0$) at $\beta > \beta_c$ in the weak coupling region, m_{5q} vanishes exponentially in N_5 and DWQCD in the finite volume works well to describe the chiral symmetry.

We think that $\beta_c < 6.0$ for the plaquette action and $\beta_c < 2.6$ for the RG improved action. At first sight this seems to contradict with the numerical data of m_{5q}, which does not show the exponential decay in N_5, for the plaquette action [14,15]. This contradicted behavior is explained as follows. In the intermediate values of N_5, the continuous eigenvalues give the main contribution of m_{5q}, so that

$$m_{5q} \sim C_{\rm cont}\exp[-N_5|\lambda_{\rm min}^{\rm cont}|], \tag{6.2}$$

while as N_5 further increases the localized eigenvalues dominates, so that

$$m_{5q} \sim C_{\rm local}\exp[-N_5|\lambda_{\rm min}^{\rm local}|], \tag{6.3}$$

where $C_{\rm cont}$ or $C_{\rm local}$ is proportional to the number of the modes near $\lambda_{\rm min}^{\rm cont}$ or $\lambda_{\rm min}^{\rm local}$, respectively. At $\beta = 6.0$ for the plaquette action, the transition between the former exponential behavior with $|\lambda_{\rm min}^{\rm cont}|$ and the latter with $|\lambda_{\rm min}^{\rm local}|$ can be seen for $N_5 = 10 \sim 40$ [14,15]. The latter behavior looks almost constant in N_5 since $\lambda_{\rm min}^{\rm local}$ is very small. At $\beta = 2.6$ for the RG improved action, on the other hand, only the former exponential behavior can be detected for $N_5 = 10 \sim 24$ [14]. This suggests that the ratio $C_{\rm local}/C_{\rm cont}$ is smaller for the RG action than for the plaquette action. Indeed it has been numerically found [29] that the number of the localized modes near $\lambda_{\rm min}^{\rm local}$ is much less for the RG action, while the number of the continuous modes near $\lambda_{\rm min}^{\rm cont}$ is similar in the two actions. (However $|\lambda_{\rm min}^{\rm local}|$ seems larger for the RG action.) We then give two predictions, which should be checked in order to test this interpretation: the exponential decay, $\exp[-N_5|\lambda_{\rm min}^{\rm local}|]$, can be seen at larger $N_5 > 60$ for the plaquette action, and the transition between the former and the latter can be seen at $N_5 = 40 \sim 60$ for the RG action.

The distribution of eigenvalues of H_W [29] suggests that even $\beta = 5.65$ for the plaquette action or $\beta = 2.2$ for the RG action is already in the weak coupling region ($\beta > \beta_c$). Indeed it seems that the transition between two exponentials has been observed in the behavior of m_{5q} at these β [14].

B. Infinite volume limit at finite N_5

In the previous subsection, we argue that m_{5q} decay exponentially in N_5 in the finite volume as long as the distribution of the continuous eigenvalues has a gap around zero. The exponential decay rate $|\lambda_{\rm min}^{\rm local}|$, however, may vanish in the infinite volume limit. Therefore, it may be the case that m_{5q} does not vanish exponentially if the infinite volume limit is taken before $N_5 \to \infty$. Moreover, the value of N_5 necessary for the suppression of the chiral symmetry breaking, m_{5q}, may increase as the volume becomes larger. This may be a disaster from the practical point of view. In this subsection, we will discuss whether the effect of the localized eigenstates to m_{5q} vanishes or not in the infinite volume limit.

In the case of the hermitian Wilson-Dirac operator, the plane-wave eigenvalues in the finite volume become the continuous spectra in the infinite volume. On the other hand, we can not predict the nature of the distribution for the localized eigenvalues in the infinite volume limit. Although the localized modes are always discrete in the finite volume, it can be continuous in the limit as will be discussed later.

We first consider the case where the number of the localized modes around zero does not grow linear in the volume. More precisely, the number of modes which satisfy $-\epsilon < \lambda < \epsilon$ with $\epsilon > 0$, denoted by $N(-\epsilon, \epsilon)$, is bounded by $c_0 \epsilon V^{c_1}$ with $c_0 > 0$ and $c_1 < 1$. In this case the contribution to m_{5q} in the infinite volume limit becomes

$$\begin{aligned} m_{5q}^{\rm localized} &\simeq \lim_{V\to\infty} \frac{1}{V} \sum_{n:{\rm localized}} e^{-|\lambda_n| N_5} < \lim_{V\to\infty} \frac{N(-\epsilon,\epsilon)}{V} + O(e^{-\epsilon N_5}) \\ &< \lim_{V\to\infty} c_0 \epsilon V^{c_1 - 1} + O(e^{-\epsilon N_5}) = O(e^{-\epsilon N_5}), \end{aligned} \tag{6.4}$$

where the sum is taken only for the localized modes. This result shows that the contribution from the localized near-zero modes vanishes in the infinite volume limit and the remaining contribution also vanishes exponentially in N_5, as long as the number of the localized near-zero modes is bounded as $N(-\epsilon, \epsilon) = c_0 \epsilon V^{c_1}$ with $c_1 < 1$. DWQCD works well in this case even in the infinite volume limit.

If the number of the localized near zero-modes is proportional to the volume; $N(-\epsilon, \epsilon) = c_0 \epsilon V$, the contribution from them remains non-zero in the infinite volume limit:

$$m_{5q}^{\rm localized} \lim_{V\to\infty} \simeq \frac{1}{V} \sum_{n:{\rm localized}} e^{-|\lambda_n| N_5} = c_0 \epsilon + O(e^{-\epsilon N_5}). \tag{6.5}$$

DWQCD does not work in this case in the infinite volume limit.

If $N(-\epsilon,\epsilon) = c_0\epsilon V$, the density of state $\rho(\lambda)$ is (almost) continuous and nonzero at $\lambda = 0$ in the infinite volume limit: $\rho(0) \neq 0$. In other words, a sum of the infinitely many δ functions from the localized modes, normalized by the volume, becomes the continuous function in the infinite volume limit. To be more precise, one should first define the integrated density of states by

$$k(\lambda) = \lim_{V\to\infty} \frac{1}{V} N(-\infty,\lambda). \tag{6.6}$$

Since $k(\lambda)$ is a monotonically increasing function in λ, its derivative, $\dfrac{dk(\lambda)}{d\lambda}$, becomes a well-defined measure. The previous statement is equivalent to that $\rho_{\rm localized}(\lambda) \equiv \dfrac{dk(\lambda)}{d\lambda}$ is continuous and non-zero at $\lambda = 0$.

Let us explain the case that $\rho_{\rm localized}(0) \neq 0$ more concretely by using a model of the eigenvalue distribution. Suppose that the discrete eigenvalues are uniformly distributed in the interval that $-1 < \lambda < 1$ and the number of modes is equal to $2cV$, so that the average interval between two successive eigenvalues becomes $1/(cV)$. In the infinite volume limit, we see that the eigenvalues are discrete but dense in the interval. If we calculate m_{5q} using this distribution, we have

$$m_{5q} = \lim_{V\to\infty} \frac{2}{V}\sum_{n=1}^{cV} e^{-\frac{n}{cV}N_5} = \lim_{V\to\infty}\frac{2}{V}\frac{e^{-\frac{N_5}{cV}}(1-e^{-N_5})}{1-e^{-\frac{N_5}{cV}}} = \frac{2c}{N_5}(1-e^{-N_5}). \tag{6.7}$$

Therefore m_{5q} does not vanish exponentially in N_5 if the infinite volume limit is first taken.

The appearance of the continuous density of states from the localized modes in the infinite volume is often observed if a kind of randomness is introduced in the interaction [33–35]. Physically the Anderson localization [36] is one of such examples. In QCD one can have infinitely many localized modes if infinitely many pairs of instanton and anti-instanton with a fixed topological charge exist. In lattice QCD, we have, in addition, very localized modes associated with the dislocations. Since the dislocations is local, the number of the discrete modes associated with them can be proportional to the volume [37,34].

Although the contribution of the localized modes to m_{5q} vanishes in the finite volume, whether it vanishes in the infinite volume limit depends on the density of states of the localized modes in the infinite volume limit. The previous numerical investigations on $\rho(0)$ [28,29] suggests $\rho(0)$ is small but non-zero at $\forall\beta \neq \infty$ in quenched QCD. Further investigations, however, are necessary in particular for the volume dependence of $\rho(0)$, to have a definite conclusion on this point whether $\rho(0)$ is non-zero or not. If $\rho(0) \neq 0$ at $\forall\beta \neq \infty$, it is interesting and important to derive the β dependence of $\rho(0)$ [28] theoretically. Note however that $\rho(0)$ should vanish in the continuum limit for $0 < M < 2$, since no zero mode appears in the free theory.

C. Phase structure with the Wilson quarks

In this subsection we consider the relation between the above interpretation for the behavior of m_{5q} in DWQCD and the parity-flavor breaking in the Wilson quark action in the quenched approximation.

We first show that the contribution of the localized modes to the parity-flavor breaking order parameter vanishes in the infinite volume limit, in the case that $N(-\epsilon,\epsilon) \le c_0\epsilon V^{c_1}$ with $c_1 < 1$. In this case we have

$$\begin{aligned}
\langle\bar\psi i\gamma_5\tau^3\psi\rangle^{\rm localized} &= \lim_{H\to 0}\lim_{V\to\infty}\frac{1}{V}\sum_{n:\rm localized}\frac{-2H}{\lambda_n^2+H^2} \\
&\le \lim_{H\to 0}\lim_{V\to\infty}\frac{1}{V}\left[\epsilon c_0 V^{c_1}\frac{-2H}{H^2}+\sum_{n:|\lambda_n|>\epsilon}\frac{-2H}{\epsilon^2+H^2}\right] \\
&\le \lim_{H\to 0}\lim_{V\to\infty} c_0\epsilon V^{c_1-1}\frac{-2}{H}+\lim_{H\to 0}\lim_{V\to\infty}\frac{1}{V}C(\epsilon)V\frac{-2H}{\epsilon^2+H^2} \\
&\le C(\epsilon)\lim_{H\to 0}\frac{-2H}{\epsilon^2+H^2}=0, \qquad (6.8)
\end{aligned}$$

where $\sum_{n:|\lambda_n|>\epsilon}1 \equiv C(\epsilon)V$. The first term vanishes in the $V\to\infty$ limit, while the second one vanishes in the $H\to 0$ limit.

We next show that the contribution of the localized modes to the parity-flavor breaking order parameter remains non-zero in the infinite volume limit, if the density of such states is non-zero at $\lambda = 0$ in the infinite volume limit: $N(-\epsilon,\epsilon) = c_0\epsilon V$. To see this, we again use the uniform distribution of eigenvalues between -1 and 1, used in the previous subsection.

$$\begin{aligned}
\langle\bar\psi i\gamma_5\tau^3\psi\rangle^{\rm localized} &= \lim_{H\to 0}\lim_{V\to\infty}\frac{2}{V}\sum_{n=1}^{cV}\frac{-2H}{(\frac{n}{cV})^2+H^2} = -4c\lim_{H\to 0}\lim_{V\to\infty}\sum_{n=1}^{cV}\frac{HcV}{n^2+(HcV)^2} \\
&= -4c\lim_{H\to 0}\lim_{V\to\infty}\left[\frac{\pi\coth(\pi HcV)}{2}-\frac{1}{2HcV}+\Delta\right] = -2c\pi, \qquad (6.9)
\end{aligned}$$

where $\lim_{H\to 0}\lim_{V\to\infty}\Delta \le \lim_{H\to 0}\pi H/(2\sqrt{1+H^2}) = 0$.

The above two considerations show that the correspondence between the failure of DWQCD in the infinite volume limit and non-zero order parameter of the parity-flavor breaking in the quenched Wilson fermion still holds even in the case that the localized modes dominate in near-zero eigenvalues. The previous investigations [28,29] suggests that the gap in the phase structure in Fig. 1 closes even at $\beta > \beta_c$ in the quenched QCD with the Wilson quark. In this case we expect that non-zero value of $\rho_{\rm localized}(0)$ in the region A is much smaller than $\rho_{\rm continuous}(0)$ in the region B, the true parity-flavor breaking phase [28], so that $\rho(0)_{\rm localized}$ may be too small to be detected by measuring the pion mass. This may be a reason why the previous investigation indicates that the region A exists at $\beta = 6.0$ for the plaquette action [27]. Finally it is noted that the region A with $\rho_{\rm localized}(0) = 0$ should always exist in full QCD, since appearances of the localized near-zero modes are suppressed by the fermion determinant, $\det D_W = \det H_W$.

D. Exceptional configurations

Finally let us comment on the exceptional configurations appeared in the quenched QCD with the Wilson-type quark action. It is sometimes observed at small quark masses in the quenched simulations that the pion propagator receives the anomalously large contribution from a particular configuration, which is called the exceptional configuration and is removed from the statistical average. In ref. [27] the several configurations, which give the W shape in the pion correlation function in t space, have been observed in the region A, where the gap(absence) of the party-flavor breaking phase is expected. It has been very difficult to understand the W shape propagator, since it means that the correlation increase as the separation t increases. Now we argue that this may be understood by the localized modes.

The pion correlation function is defined by

$$\begin{aligned}\langle \bar\psi i\gamma_5\tau^a\psi(x)\cdot\bar\psi i\gamma_5\tau^a\psi(y)\rangle &= \left\langle x\left|\frac{1}{H_w}\right|y\right\rangle\left\langle y\left|\frac{1}{H_w}\right|x\right\rangle \\ &= \sum_n \langle x|\lambda_n\rangle\frac{1}{\lambda_n}\langle\lambda_n|y\rangle\sum_l\langle y|\lambda_l\rangle\frac{1}{\lambda_l}\langle\lambda_l|x\rangle \\ &= \sum_{n,l}\frac{1}{\lambda_n\lambda_l}\phi_n(x)\phi_l(x)^\dagger\phi_l(y)\phi_n(y)^\dagger, \end{aligned} \tag{6.10}$$

where $\phi_n(x)=\langle x|\lambda_n\rangle$ etc. If some localized eigenstate has a very small eigenvalue λ_n on some configuration, the contribution of the mode becomes very large:

$$\langle\bar\psi i\gamma_5\tau^a\psi(x)\cdot\bar\psi i\gamma_5\tau^a\psi(y) = \frac{1}{\lambda_n^2}\phi_n(x)\phi_n(x)^\dagger\phi_n(y)\phi_n(y)^\dagger + \text{other contributions.} \tag{6.11}$$

This is the exceptional configuration. Furthermore suppose the eigenstate $\phi_n(\vec{x},t)$ is localized at $x=(\vec{x}_n,t_n)$. Summing over $\vec{x}$ and $\vec{y}$, and taking $x_0=t$ and $y_0=0$, the time correlation is dominated by

$$C(t)\equiv\sum_{\vec{x},\vec{y}}\frac{1}{\lambda_n^2}\phi_n(\vec{x},t)\phi_n(\vec{x},t)^\dagger\phi_n(\vec{y},0)\phi_n(\vec{y},0)^\dagger, \tag{6.12}$$

and $C(t)$ has a peak around $t=t_n$. Together with an enhance factor $1/\lambda_n^2$ for small eigenvalues this explains the W shape behavior of the pion propagator.

VII. CONCLUSIONS

In this paper we first derive the formula for the chiral symmetry breaking term m_{5q} in DWQCD in terms of the (modified) hermitian Wilson-Dirac operator in 4 dimensions. Using several simplifications and approximations we explicitly write down the formula of m_{5q} in terms the eigenvalues only.

The important observation is that there are two different types for the eigenvalues of the the hermitian Wilson-Dirac operator. One is the continuous one, which corresponds to the plane-wave in the free theory, the other is the localized one, associated with the instanton or dislocations. We argue that the effect of the latter one to the physical observable such as m_{5q} or the parity-flavor breaking order parameter vanishes in the infinite volume limit, unless the number of near zero localized modes increases linearly in the volume. On the other hand, the effect remains non-zero but small in the infinite volume limit, if it linearly increases.

The message of this paper is that domain-wall fermion or overlap Dirac fermion should work well to describe the vector-like chiral symmetry at weak coupling, $\beta>\beta_c$, where the gap in the continuous spectra opens, as long as the volume is finite. The small eigenvalues appeared in the numerical simulation seem to belong to the localized one, hence their contribution vanishes in the $N_5 \to \infty$ limit in the finite volume. Whether domain-wall/overlap fermions are successful or not in the infinite volume, however, depends crucially on the distribution of the localized eigenvalues in the limit.

In practice one can make such small eigenvalues appeared in the finite volume larger by hand, without changing physical observables [38,39]. This is a little costly. If the effect of chiral symmetry breaking is small enough or no dependence of observables on N_5 is detected, one may instead perform simulations at large but numerically affordable value of N_5. Except the quantities very sensitive to the small eigenvalues such as m_{5q}, N_5 dependences are expected to be rather week in general. This expectation is indeed true in the case of the quantum hall effect [40].

ACKNOWLEDGMENTS

We thank Profs. Y. Kikukawa, T. Izubuchi and K. I. Nagai for useful discussion. S.A would like to thank Profs. M. Lüscher, H. Neuberger and S. Nakamura for informative discussion, and Y.T would like to thank Prof. T. Onogi for valuable discussion. This work is supported in part by Grants-in-Aid of the Ministry of Education (Nos. 12304011,12640253, 13135204).

APPENDIX A: PLANE WAVE

The eigenvector (4.25) of the free Hamiltonian (4.18) is given by combining positive and negative energy eigenvector $u_\alpha(p,\tau)$ and $v_\alpha(p,\tau)$

$$(H')_{\alpha\beta} v_\beta(p,\tau) = -|\lambda'(p)| v_\alpha(p,\tau), \tag{A1}$$

$$(H')_{\alpha\beta} u_\beta(p,\tau) = |\lambda'(p)| u_\alpha(p,\tau), \tag{A2}$$

where α, β are spinor indexes and τ runs $\tau = 1, 2$. We adopt the following combination in this paper;

$$U_\alpha(p,1) = v_\alpha(p,1),\ U_\alpha(p,2) = v_\alpha(p,2),\ U_\alpha(p,3) = u_\alpha(p,1),\ U_\alpha(p,4) = u_\alpha(p,2). \tag{A3}$$

u and v are given as follows with Pauli matrix $\vec{\sigma}$ and two dimensional basis vector $\xi(\tau)$,

$$v_\alpha(p,\tau) = \sqrt{\frac{|\lambda'(p)| - B(p)}{2|\lambda'(p)|}} \begin{pmatrix} \xi(\tau) \\ \frac{-i\sigma^\dagger_\mu A_\mu(p)}{B(p)-|\lambda'(p)|}\xi(\tau) \end{pmatrix}_\alpha, \tag{A4}$$

$$u_\alpha(p,\tau) = \sqrt{\frac{|\lambda'(p)| - B(p)}{2|\lambda'(p)|}} \begin{pmatrix} \frac{-i\sigma_\mu A_\mu(p)}{B(p)-|\lambda'(p)|}\xi(\tau) \\ \xi(\tau) \end{pmatrix}_\alpha, \tag{A5}$$

$$\sigma_\mu = (1, -i\vec{\sigma}), \tag{A6}$$

$$\xi(1) = \begin{pmatrix} 1 \\ 0 \end{pmatrix},\ \xi(2) = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \tag{A7}$$

$U_\alpha(p,s)$ satisfies the orthogonal and completeness condition

$$\sum_\alpha U_\alpha^\dagger(p,s)U_\alpha(p,t) = \delta_{s,t}, \tag{A8}$$

$$\sum_s U_\alpha^\dagger(p,s)U_\beta(p,s) = \delta_{\alpha,\beta}. \tag{A9}$$

APPENDIX B: ANOMALOUS QUARK MASS WITH COMPLETELY LOCALIZED AND PLANE-WAVE EIGENVECTORS

We consider a system with N_l completely localized eigenvectors and plane-wave functions with N_p degrees of freedom for momentum p.

We expand the two Green functions in m_{5q} with the complete set in eq.(4.35).

$$\begin{aligned}
X(t) &= \sum_{\vec{x}} \langle J_{5q}(x)P(y)\rangle \\
&= -\frac{a_5}{a^{10}} \sum_{\vec{x},\alpha,a}\sum_{b,\beta} \left\langle I \left| \frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}} D_{N_5}^{-1} \right| J \right\rangle \left\langle J \left| \left(D_{N_5}^{-1}\right)^\dagger \frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}} \right| I \right\rangle \\
&= -\frac{a_5}{a^{10}} \sum_{\vec{x},\alpha,a}\sum_{b,\beta} \Bigg(\sum_{n,m=1}^{N_l} \langle I|n\rangle \left\langle n \left| \frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}} D_{N_5}^{-1} \right| J \right\rangle \\
&\quad \times \left\langle J \left| \left(D_{N_5}^{-1}\right)^\dagger \frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}} \right| m \right\rangle \langle m|I\rangle \\
&\quad + \sum_{p,s,c}\sum_{k,t,d} \langle I|p,s,c\rangle \left\langle p,s,c \left| \frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}} D_{N_5}^{-1} \right| J \right\rangle \\
&\quad \times \left\langle J \left| \left(D_{N_5}^{-1}\right)^\dagger \frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}} \right| k,t,d \right\rangle \langle k,t,d|I\rangle \Bigg),
\end{aligned} \tag{B1}$$

where $I=(x,\alpha,a)$, $J=(y,\beta,b)$ and we use a relation $\langle I|n\rangle\langle k,t,d|I\rangle = 0$.

Contribution from the localized eigenvector is given as follows

$$\begin{aligned}
X_l(t) &= -\frac{a_5}{a^{10}} \sum_{\vec{x},\alpha,a}\sum_{b,\beta}\sum_{n,m=1}^{N_l} \langle I|n\rangle\langle m|I\rangle \\
&\quad \times \left\langle n \left| \frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}} D_{N_5}^{-1} \right| J \right\rangle \left\langle J \left| \left(D_{N_5}^{-1}\right)^\dagger \frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}} \right| m \right\rangle \\
&= -\frac{a_5}{a^{10}} \sum_{n=1}^{N_l} \delta_{t,x_n^0} \left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda_n')} \right)^2 \sum_{b,\beta} \left\langle n \left| D_{N_5}^{-1} \right| y,\beta,b \right\rangle \left\langle y,\beta,b \left| \left(D_{N_5}^{-1}\right)^\dagger \right| n \right\rangle,
\end{aligned} \tag{B2}$$

where we use a relation

$$\langle I|n\rangle = \langle x,\alpha,a|n\rangle = \delta_{\alpha,\alpha_n}\delta_{a,a_n}\delta_{x,x_n}, \tag{B3}$$

$$\langle I|n\rangle\langle m|I\rangle = \delta_{m,n}\delta_{\alpha,\alpha_n}\delta_{a,a_n}\delta_{x,x_n}. \tag{B4}$$

By inserting the identity operator in (x, α, a) space

$$1 = \sum_{x,\alpha,a} |x, \alpha, a\rangle \langle x, \alpha, a| \tag{B5}$$

we have

$$\begin{aligned}
X_l(t) &= -\frac{a_5}{a^{10}} \sum_{n=1}^{N_l} \delta_{t,x_n^0} \left(\frac{1}{2\cosh \frac{N_5}{2} a_5 \widetilde{\lambda}(\lambda'_n)} \right)^2 \\
&\quad \times \sum_{b,\beta} \sum_{z,\gamma,c} \sum_{w,\delta,d} \langle n|z,\gamma,c\rangle \left\langle z,\gamma,c \left| D_{N_5}^{-1} \right| y,\beta,b \right\rangle \left\langle y,\beta,b \left| \left(D_{N_5}^{-1}\right)^\dagger \right| w,\delta,d \right\rangle \langle w,\delta,d|n\rangle \\
&= -\frac{a_5}{a^{10}} \sum_{n=1}^{N_l} \delta_{t,x_n^0} \left(\frac{1}{2\cosh \frac{N_5}{2} a_5 \widetilde{\lambda}(\lambda'_n)} \right)^2 \\
&\quad \times \sum_{b,\beta} \left\langle x_n,\alpha_n,a_n \left| D_{N_5}^{-1} \right| y,\beta,b \right\rangle \left\langle y,\beta,b \left| \left(D_{N_5}^{-1}\right)^\dagger \right| x_n,\alpha_n,a_n \right\rangle \\
&= -\frac{a_5}{a^{10}} \frac{1}{4N_c N_x} \sum_{n=1}^{N_l} \left(\frac{1}{2\cosh \frac{N_5}{2} a_5 \widetilde{\lambda}(\lambda'_n)} \right)^2 \\
&\quad \times \sum_{\vec{x},\alpha,a} \sum_{b,\beta} \left\langle \vec{x},t;\alpha,a \left| D_{N_5}^{-1} \right| y,\beta,b \right\rangle \left\langle y,\beta,b \left| \left(D_{N_5}^{-1}\right)^\dagger \right| \vec{x},t;\alpha,a \right\rangle \\
&= \frac{1}{a_5} Y(t) \frac{1}{4N_c N_x} \sum_{n=1}^{N_l} \left(\frac{1}{2\cosh \frac{N_5}{2} a_5 \widetilde{\lambda}(\lambda'_n)} \right)^2,
\end{aligned} \tag{B6}$$

where we assume that the center of the localized solution (x_n, α_n, a_n) is distributed uniformly in the (x, α, a) space and we can take average over (x, α, a) when enough number of configurations are summed in the simulation.

For the contribution from the continuous eigenvalues we take $t \to \infty$ limit to proceed,

$$\begin{aligned}
X_c(t)|_{t\to\infty} &= -\frac{a_5}{a^{10}} \sum_{\vec{x},\alpha,a} \sum_{b,\beta} \sum_{p,s} \sum_{k,s'} \frac{1}{N_p} U_\alpha(p,s) U_\alpha^\dagger(k,t) e^{i(p-k)x} \\
&\quad \times \left\langle p,s,a \left| \frac{1}{2\cosh \frac{N_5}{2} a_5 \widetilde{H}} D_{N_5}^{-1} \right| y,\beta,b \right\rangle \left\langle y,\beta,b \left| \left(D_{N_5}^{-1}\right)^\dagger \frac{1}{2\cosh \frac{N_5}{2} a_5 \widetilde{H}} \right| k,s',a \right\rangle \\
&= -\frac{a_5}{a^{10}} n'_{xyz} \frac{1}{N_p} \sum_{p,s,a} \left(\frac{1}{2\cosh \frac{N_5}{2} a_5 \widetilde{\lambda}(p)} \right)^2 \\
&\quad \times \sum_{\beta,b} \left\langle p,s,a \left| D_{N_5}^{-1} \right| y,\beta,b \right\rangle \left\langle y,\beta,b \left| \left(D_{N_5}^{-1}\right)^\dagger \right| p,s,a \right\rangle,
\end{aligned} \tag{B7}$$

where we use following relations

$$\langle x,\alpha,a|p,s,b\rangle = \frac{1}{\sqrt{N_p}} U_\alpha(p,s) e^{ipx} \delta_{a,b}, \tag{B8}$$

$$\sum_{\vec{x}} e^{i(\vec{p}-\vec{k})\vec{x}} = n'_{xyz} \delta_{\vec{p},\vec{k}}, \tag{B9}$$

$$\lim_{t\to\infty} e^{it(p^0-k^0)} = \delta_{p^0,k^0}, \tag{B10}$$

$$\sum_\alpha U^\dagger_\alpha(p,s)U_\alpha(p,s') = \delta_{s,s'}. \tag{B11}$$

n'_{xyz} is a volume of $\widetilde{\mathcal{S}}$ in xyz space. By inserting the complete set (4.35) of eigenstates we have

$$\begin{aligned}
X_c(t)|_{t\to\infty} = &-\frac{a_5}{a^{10}}n'_{xyz}\frac{1}{N_p}\sum_{p,s,a}\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(p)}\right)^2 \\
&\times\sum_{\beta,b}\Biggl(\sum_{n,m=1}^{N_l}\left\langle p,s,a\left|D^{-1}_{N_5}\right|n\right\rangle\langle n|y,\beta,b\rangle\langle y,\beta,b|m\rangle\left\langle m\left|\left(D^{-1}_{N_5}\right)^\dagger\right|p,s,a\right\rangle \\
&+\sum_{k,s',c}\left\langle p,s,a\left|D^{-1}_{N_5}\right|k,s',c\right\rangle\langle k,s',c|y,\beta,b\rangle \\
&\times\sum_{q,s'',d}\langle y,\beta,b|q,s'',d\rangle\left\langle q,s'',d\left|\left(D^{-1}_{N_5}\right)^\dagger\right|p,s,a\right\rangle\Biggr) \\
=&-\frac{a_5}{a^{10}}n'_{xyz}\frac{1}{N_p}\sum_p\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(p)}\right)^2 \\
&\times\Biggl(\sum_{a,s}\sum_{n=1}^{N_l}\delta_{y,x_n}\left\langle p,s,a\left|D^{-1}_{N_5}\right|n\right\rangle\left\langle n\left|\left(D^{-1}_{N_5}\right)^\dagger\right|p,s,a\right\rangle \\
&+\frac{N_c}{N_p}\sum_{s,s'}2a\left(\frac{-i\gamma_\mu C_\mu(p)+E(p)}{C^2+E^2}\right)_{s,s'}2a\left(\frac{i\gamma_\mu C_\mu(p)+E(p)}{C^2+E^2}\right)_{s',s}\Biggr) \\
=&-\frac{a_5}{a^8}n'_{xyz}\frac{1}{N_p}\sum_p\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(p)}\right)^2F(p),
\end{aligned} \tag{B12}$$

where $F(p)$ is defined as

$$F(p) = F_l(p)+F_c(p), \tag{B13}$$

$$F_l(p) = \sum_{n=1}^{N_l}\delta_{y,x_n}\sum_{s,a}\frac{1}{a^2}\left\langle p,s,a\left|D^{-1}_{N_5}\right|n\right\rangle\left\langle n\left|\left(D^{-1}_{N_5}\right)^\dagger\right|p,s,a\right\rangle, \tag{B14}$$

$$F_c(p) = \frac{4N_c}{N_p}\left(\frac{4}{C^2(p)+E^2(p)}\right). \tag{B15}$$

We can show that $F_l(p)$ vanishes as follows. We extend the orthogonality assumption of the eigenvectors $\langle n|p,s,a\rangle = 0$ to the case with γ_5; $\langle n|\gamma_5|p,s,a\rangle = 0$. This is plausible since the eigenfunction $\psi_n(x) = \langle x|n\rangle$ is localized in space-time and the overlap is suppressed even if we multiply γ_5. Then by using the explicit form (3.5), the matrix element of the truncated overlap Dirac operator D_{N_5} turns out to be block diagonal, in which only the matrix elements $\langle n|D_{N_5}|m\rangle$ and $\langle p,s,a|D_{N_5}|k,t,b\rangle$ are non-zero and the off-diagonal part $\langle p,s,a|D_{N_5}|n\rangle$ becomes zero. This block diagonal property is kept even if we take inversion and the off-diagonal part becomes zero

$$\left\langle p,s,a\left|D^{-1}_{N_5}\right|n\right\rangle = 0. \tag{B16}$$

It is noted that in the free theory the relation $D_W^\dagger D_W = D_W D_W^\dagger$ is satisfied and $\left\langle n \left| D_{N_5}^{-1} \right| m \right\rangle = 0$ is shown exactly for non-degenerate $|\lambda'_n| \neq |\lambda'_m|$.

The pion propagator in the denominator of m_{5q} is expanded similarly with eigenstates as

$$\begin{aligned}
Y(t) &= \sum_{\vec{x}} \langle P(x)P(y) \rangle \\
&= -\frac{a_5^2}{a^{10}} \sum_{\vec{x},a,\alpha} \sum_{b,\beta} \left\langle I \left| D_{N_5}^{-1} \right| J \right\rangle \left\langle J \left| (D_{N_5}^{-1})^\dagger \right| I \right\rangle \\
&= -\frac{a_5^2}{a^{10}} \sum_{\vec{x},a,\alpha} \sum_{b,\beta} \Bigg(\sum_{n,m=1}^{N_l} \langle I|n \rangle \left\langle n \left| D_{N_5}^{-1} \right| J \right\rangle \left\langle J \left| (D_{N_5}^{-1})^\dagger \right| m \right\rangle \langle m|I \rangle \\
&\quad + \sum_{p,s,c} \sum_{k,s',d} \langle I|p,s,c \rangle \left\langle p,s,c \left| D_{N_5}^{-1} \right| J \right\rangle \left\langle J \left| (D_{N_5}^{-1})^\dagger \right| k,s',d \right\rangle \langle k,s',d|I \rangle \Bigg).
\end{aligned} \tag{B17}$$

A contribution from localized eigenvector is given by

$$\begin{aligned}
Y_l(t) &= -\frac{a_5^2}{a^{10}} \sum_{n=1}^{N_l} \delta_{t,x_n^0} \sum_{b,\beta} \left\langle n \left| D_{N_5}^{-1} \right| y,\beta,b \right\rangle \left\langle y,\beta,b \left| (D_{N_5}^{-1})^\dagger \right| n \right\rangle \\
&= -\frac{a_5^2}{a^{10}} \sum_{n=1}^{N_l} \delta_{t,x_n^0} \sum_{b,\beta} \sum_{z,\gamma,c} \langle n|z,\gamma,c \rangle \left\langle z,\gamma,c \left| D_{N_5}^{-1} \right| y,\beta,b \right\rangle \\
&\quad \times \sum_{w,\delta,d} \left\langle y,\beta,b \left| (D_{N_5}^{-1})^\dagger \right| w,\delta,d \right\rangle \langle w,\delta,d|n \rangle \\
&= -\frac{a_5^2}{a^{10}} \sum_{n=1}^{N_l} \delta_{t,x_n^0} \sum_{b,\beta} \left\langle x_n,\alpha_n,a_n \left| D_{N_5}^{-1} \right| y,\beta,b \right\rangle \left\langle y,\beta,b \left| (D_{N_5}^{-1})^\dagger \right| x_n,\alpha_n,a_n \right\rangle \\
&= -\frac{a_5^2}{a^{10}} \sum_{n=1}^{N_l} \frac{1}{4N_c N_x} \sum_{\vec{x},\alpha,a} \sum_{\beta,b} \left\langle \vec{x},t,\alpha,a \left| D_{N_5}^{-1} \right| y,\beta,b \right\rangle \left\langle y,\beta,b \left| (D_{N_5}^{-1})^\dagger \right| \vec{x},t,\alpha,a \right\rangle \\
&= \frac{N_l}{4N_c N_x} Y(t).
\end{aligned} \tag{B18}$$

Taking $t \to \infty$ limit a contribution from the continuous sector becomes

$$\begin{aligned}
Y_c(t)|_{t\to\infty} &= -\frac{a_5^2}{a^{10}} \sum_{\vec{x},a,\alpha} \sum_{b,\beta} \sum_{p,s} \sum_{k,s'} \frac{1}{N_p} U_\alpha(p,s) U_\alpha^\dagger(k,s') e^{i(p-k)x} \\
&\quad \times \left\langle p,s,a \left| D_{N_5}^{-1} \right| y,\beta,b \right\rangle \left\langle y,\beta,b \left| (D_{N_5}^{-1})^\dagger \right| k,s',a \right\rangle \\
&= -\frac{a_5^2}{a^{10}} n'_{xyz} \frac{1}{N_p} \sum_{p,s,a} \sum_{b,\beta} \left\langle p,s,a \left| D_{N_5}^{-1} \right| y,\beta,b \right\rangle \left\langle y,\beta,b \left| (D_{N_5}^{-1})^\dagger \right| p,s,a \right\rangle \\
&= -\frac{a_5^2}{a^{10}} n'_{xyz} \frac{1}{N_p} \sum_{p,s,a} \\
&\quad \times \sum_{b,\beta} \Bigg(\sum_{n,m=1}^{N_l} \left\langle p,s,a \left| D_{N_5}^{-1} \right| n \right\rangle \langle n|y,\beta,b \rangle \langle y,\beta,b|m \rangle \left\langle m \left| (D_{N_5}^{-1})^\dagger \right| p,s,a \right\rangle
\end{aligned}$$

$$
\begin{aligned}
&+\sum_{k,s',c}\left\langle p,s,a\left|D_{N_5}^{-1}\right|k,s',c\right\rangle\langle k,s',c|y,\beta,b\rangle\\
&\quad\times\sum_{q,s'',d}\langle y,\beta,b|q,s'',d\rangle\left\langle q,s'',d\left|(D_{N_5}^{-1})^{\dagger}\right|p,s,a\right\rangle\Bigg)\\
=&-\frac{a_5^2}{a^{10}}n'_{xyz}\frac{1}{N_p}\sum_p F(p).
\end{aligned}
\tag{B19}
$$

As a consequence we have a relation

$$
Y(t\to\infty)=-\frac{4N_cN_x}{4N_cN_x-N_l}\frac{a_5^2}{a^{10}}n'_{xyz}\frac{1}{N_p}\sum_p F(p).
\tag{B20}
$$

Now we need to know behavior of $F(p)$ as a function of p to further proceed. We can directly see that the function $F(p)=F_c(p)$ is almost constant for all the range of p as in Fig. 2. We assume that $F(p)$ is independent of p.

By adopting this assumption the anomalous quark mass becomes

$$
\begin{aligned}
m_{5q}&=\frac{\lim_{t\to\infty}X(t)}{\lim_{t\to\infty}Y(t)}\\
&=\frac{1}{a_5}\frac{1}{4N_cN_x}\sum_n\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda'_n)}\right)^2\\
&\quad-\frac{1}{Y(t\to\infty)}\frac{a_5}{a^{10}}n'_{xyz}\frac{1}{N_p}\sum_p\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(p)}\right)^2F(p)\\
&=\frac{1}{a_5}\frac{1}{4N_cN_x}\sum_n\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda'_n)}\right)^2\\
&\quad+\frac{4N_cN_x-N_l}{4N_cN_x}\frac{1}{a_5}\frac{1}{\sum_k F(k)}\sum_p\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(p)}\right)^2F(p)\\
&=\frac{1}{a_5}\frac{1}{4N_cN_x}\left(\sum_n\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda'_n)}\right)^2+4N_c\sum_p\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(p)}\right)^2\right)\\
&=\frac{1}{a_5}\frac{1}{4N_cN_x}\left(\sum_n\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda'_n)}\right)^2+\sum_n\rho_n\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}_n}\right)^2\right),
\end{aligned}
\tag{B21}
$$

where ρ_n is a number of degeneracy in $\widetilde{\lambda}_n$,

$$
\rho_n=\sum_{p,\alpha,a}\delta_{\widetilde{\lambda}_n,\widetilde{\lambda}(p)}.
\tag{B22}
$$

APPENDIX C: ANOMALOUS QUARK MASS WITH PARTIALLY LOCALIZED AND PLANE-WAVE EIGENVECTORS

We consider a system with N_l subspaces $\mathcal{S}_n$ $(n = 1, \cdots, N_l)$, which is spanned by partially localized eigenvectors $\psi_n^{(i)}$. We assume that each subspace $\mathcal{S}_n$ has no overlap and its volume is h_n. Remaining subspace $\widetilde{\mathcal{S}}$ of the system is spanned by plane-wave eigenvectors as in the previous subsection. If we set the size of $\widetilde{\mathcal{S}}$ to be $4N_cN_p$ the total degrees of freedom of the system becomes

$$4N_cN_x = \sum_{n=1}^{N_l} h_n + 4N_cN_p. \tag{C1}$$

The partially localized eigenvector is given by

$$H' |n, i\rangle = \lambda_n^{(i)} |n, i\rangle \,, \tag{C2}$$

$$\langle I|n, i\rangle = \frac{1}{\sqrt{h_n}} \psi_n^{(i)}(I), \tag{C3}$$

where the function $\psi_n^{(i)}(I)$ is non-zero within $\mathcal{S}_n$. We have a relation that

$$\psi_n^{(i)\dagger}(I)\psi_m^{(j)}(I) = 0 \tag{C4}$$

for $n \neq m$ for any i, j. The orthogonality is given by

$$\sum_I \psi_n^{(i)\dagger}(I)\psi_m^{(j)}(I) = h_n\delta_{n,m}\delta_{i,j}. \tag{C5}$$

The plane-wave eigenvector is the same as in the previous section (4.25) with a constraint

$$\langle n, i|I\rangle \langle I|p, s, b\rangle = 0. \tag{C6}$$

The completeness condition of this system is given by

$$1 = \sum_{n=1}^{N_l} \sum_{i=1}^{h_n} |n, i\rangle \langle n, i| + \sum_{l,s} \sum_p \sum_a |p, l, s, a\rangle \langle p, l, s, a| \tag{C7}$$

with

$$\sum_{n=1}^{N_l} \sum_{i=1}^{h_n} \frac{1}{h_n} \psi_n^{(i)}(I)\psi_n^{(i)\dagger}(J) = \delta_{I,J}. \tag{C8}$$

We expand the two Green functions in m_{5q} with the complete set in the above.

$$\begin{aligned} X(t) &= \sum_{\vec{x}} \langle J_{5q}(x)P(y)\rangle \\ &= -\frac{a_5}{a^{10}} \sum_{\vec{x},\alpha,a} \sum_{b,\beta} \left\langle I \left| \frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}} D_{N_5}^{-1} \right| J \right\rangle \left\langle J \left| \left(D_{N_5}^{-1}\right)^\dagger \frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}} \right| I \right\rangle \end{aligned}$$

$$
\begin{aligned}
&= -\frac{a_5}{a^{10}} \sum_{\vec{x},\alpha,a}\sum_{b,\beta}\Bigg(\sum_{n,m=1}^{N_l}\sum_{i,j=1}^{h_n} \langle x,\alpha,a|n,i\rangle \left\langle n,i\left|\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}}D_{N_5}^{-1}\right|J\right\rangle \\
&\quad\times \left\langle J\left|\left(D_{N_5}^{-1}\right)^{\dagger} \frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}}\right|m,j\right\rangle \langle m,j|x,\alpha,a\rangle \\
&\quad+ \sum_{p,s,c}\sum_{k,t,d}\langle x,\alpha,a|p,s,c\rangle \left\langle p,s,c\left|\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}}D_{N_5}^{-1}\right|J\right\rangle \\
&\quad\times \left\langle J\left|\left(D_{N_5}^{-1}\right)^{\dagger} \frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}}\right|k,t,d\right\rangle \langle k,t,d|x,\alpha,a\rangle\Bigg).
\end{aligned}
\tag{C9}
$$

The contribution X_c from the continuous eigenvalues is the same as in the previous subsection except for the definition of $F(p)$

$$F(p) = F_l(p) + F_c(p), \tag{C10}$$

$$F_l(p) = \sum_{n=1}^{N_l}\sum_{i,j=1}^{h_n}\sum_{p,s,a}\left\langle p,s,a\left|D_{N_5}^{-1}\right|n,i\right\rangle \frac{1}{h_n}\psi_n^{(i)\dagger}(J)\psi_n^{(j)}(J)\left\langle n,j\left|\left(D_{N_5}^{-1}\right)^{\dagger}\right|p,s,a\right\rangle, \tag{C11}$$

$$F_c(p) = \frac{4N_c}{N_p}\left(\frac{4}{C^2(p)+E^2(p)}\right). \tag{C12}$$

Here we use the block diagonal condition

$$\left\langle p,s,a\left|D_{N_5}^{-1}\right|n,i\right\rangle = 0 \tag{C13}$$

by assuming the γ_5 orthogonality $\langle n,i\,|\gamma_5|\,p,s,a\rangle = 0$. X_c is written in terms of the function $F(p) = F_c(p)$.

Contribution from the localized mode becomes

$$
\begin{aligned}
X_l(t) &= -\frac{a_5}{a^{10}}\sum_{b,\beta}\sum_{n=1}^{N_l}\sum_{i,j=1}^{h_n}\frac{1}{h_n}f_n^{i,j}(t) \\
&\quad\times\left\langle n,i\left|\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}}D_{N_5}^{-1}\right|J\right\rangle\left\langle J\left|\left(D_{N_5}^{-1}\right)^{\dagger}\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{H}}\right|n,j\right\rangle \\
&= -\frac{a_5}{a^{10}}\sum_{n=1}^{N_l}\sum_{i,j=1}^{h_n}\frac{1}{h_n}f_n^{i,j}(t)\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda_n^{\prime(i)})}\right)\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda_n^{\prime(j)})}\right) \\
&\quad\times\sum_{b,\beta}\sum_{z,\gamma,c}\sum_{w,\delta,d}\langle n,i|z,\gamma,c\rangle\left\langle z,\gamma,c\left|D_{N_5}^{-1}\right|J\right\rangle\left\langle J\left|\left(D_{N_5}^{-1}\right)^{\dagger}\right|w,\delta,d\right\rangle\langle w,\delta,d|n,j\rangle \\
&= -\frac{a_5}{a^{10}}\sum_{n=1}^{N_l}\sum_{i,j=1}^{h_n}\frac{1}{h_n}f_n^{i,j}(t)\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda_n^{\prime(i)})}\right)\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda_n^{\prime(j)})}\right) \\
&\quad\times\sum_{b,\beta}\sum_{z,\gamma,c}\sum_{w,\delta,d}\frac{1}{h_n}\psi_n^{(i)\dagger}(z,\gamma,c)\left\langle z,\gamma,c\left|D_{N_5}^{-1}\right|J\right\rangle\left\langle J\left|\left(D_{N_5}^{-1}\right)^{\dagger}\right|w,\delta,d\right\rangle\psi_n^{(j)}(w,\delta,d),
\end{aligned}
\tag{C14}
$$

where we assumed orthogonality for m and n, $f_n^{i,j}$ is defined as

$$\sum_{\vec{x}}\sum_{\alpha,a}\psi_n^{(i)}(I)\psi_m^{(j)\dagger}(I)=\delta_{n,m}f_n^{i,j}(t). \tag{C15}$$

$f_n^{i,j}(t)$ is non-zero when $t\in\mathcal{S}_n$. Here we have a comment. It is plausible to assume that the eigenfunction $\psi_n^{(i)}$ tends to be plane but rapidly oscillating inside $\mathcal{S}_n$ for large i. By using this fact we can see that the above $f_n^{i,j}$ is suppressed for different i, j even if the summation over t is not taken.

We need more information on the propagator multiplied with eigenvectors

$$\frac{1}{h_n}\sum_I\sum_K\psi_n^{(i)\dagger}(I)\left\langle I\left|D_{N_5}^{-1}\right|J\right\rangle\left\langle J\left|\left(D_{N_5}^{-1}\right)^\dagger\right|K\right\rangle\psi_n^{(j)}(K) \tag{C16}$$

to further discuss the detailed property of X_l. We start by adopting an assumption that the propagator

$$f(I,K)=\left\langle I\left|D_{N_5}^{-1}\right|J\right\rangle\left\langle J\left|\left(D_{N_5}^{-1}\right)^\dagger\right|K\right\rangle \tag{C17}$$

is a slowly varying function of I and K. Then we investigate the behavior of (C16) for two typical forms of eigenfunctions. The eigenfunctions are classified into two types, according to the behavior of the following integral of a single function

$$C=\sum_I\psi_n^{(i)}(I). \tag{C18}$$

(i) Eigenfunction $\psi_n^{(i)}$ with $C\neq 0$, which may be typical for the lowest mode $i=1$. It seems to be plausible [19,31] to approximate this lowest mode with the exponential form. For simplicity we consider a one-dimensional case

$$\psi(x)=\frac{1}{\sqrt{\delta}}e^{-|x-x_0|/\delta}, \tag{C19}$$

where δ is a width and x_0 is a center of the eigenfunction. Since the integral $\int dx\psi(x)$ gives non-zero value, a multiplication with a smooth function $f(x,y)$ produces a enhance factor 4δ, which is proportional to a width of the eigenfunction:

$$\begin{aligned}&\frac{1}{\delta}\int_{-\infty}^{\infty}dxdye^{-|x-x_0|/\delta}f(x,y)e^{-|y-x_0|/\delta}\\&=\frac{1}{\delta}\int_{-\infty}^{\infty}dzdwe^{-|z|/\delta}f(z+x_0,w+x_0)e^{-|w|/\delta}\\&=\frac{1}{\delta}\int_{-\infty}^{\infty}dzdwe^{-|z|/\delta}f(x_0,x_0)e^{-|w|/\delta}+\mathcal{O}(\partial f(x_0,x_0))=(4\delta)\ f(x_0,x_0).\end{aligned} \tag{C20}$$

The extension to four-dimensions is given by

$$\psi_n^{(1)}(x)=\frac{1}{\delta^2}e^{-\sum_i|x_i-(x_n)_i|/\delta} \tag{C21}$$

and (C16) becomes

$$\begin{aligned}
&\frac{1}{h_n}\sum_I\sum_K \psi_n^{(i)\dagger}(I)\left\langle I\left|D_{N_5}^{-1}\right|J\right\rangle\left\langle J\left|\left(D_{N_5}^{-1}\right)^\dagger\right|K\right\rangle\psi_n^{(j)}(K)\\
&=4^4h_n\delta_{i,j}\left\langle I_n\left|D_{N_5}^{-1}\right|J\right\rangle\left\langle J\left|\left(D_{N_5}^{-1}\right)^\dagger\right|I_n\right\rangle,
\end{aligned}\tag{C22}$$

where I_n is a peak of the exponential in $\psi_n^{(i)}(I)$ and h_n is given by $h_n=\delta^4$. Here we use a property that the eigenfunction $\psi_n^{(i)}$ with higher $i>1$ tends to be oscillating and a single summation $\sum_I\psi_n^{(i)}(I)$ is suppressed. Together with the suppression of $f_n^{i,j}$ for $i\neq j$ we have a factor $\delta_{i,j}$ in the above.

The term X_l of the numerator for the contribution from the lowest eigenvector $i=1$ becomes

$$\begin{aligned}
X_l(t)&=-\frac{a_5}{a^{10}}\sum_{n=1}^{N_l}\sum_{i=1}^{h_n}\delta_{i,1}4^4f_n^{i,i}(t)\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda_n^{'(i)})}\right)^2\sum_{b,\beta}\left\langle I_n\left|D_{N_5}^{-1}\right|J\right\rangle\left\langle J\left|\left(D_{N_5}^{-1}\right)^\dagger\right|I_n\right\rangle\\
&=-\frac{a_5}{a^{10}}\frac{1}{4N_cn_{xyz}}\sum_{n=1}^{N_l}\frac{4^4h_n}{h_{n,t}}\delta_{t\in\mathcal{S}_n}\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda_n^{'(1)})}\right)^2\\
&\quad\times\sum_{\vec{x}}\sum_{\alpha,a}\sum_{b,\beta}\left\langle\vec{x},t,\alpha,a\left|D_{N_5}^{-1}\right|y,\beta,b\right\rangle\left\langle y,\beta,b\left|\left(D_{N_5}^{-1}\right)^\dagger\right|\vec{x},t,\alpha,a\right\rangle,\\
&=\frac{1}{a_5}Y(t)\frac{4^4h_n}{4N_cN_x}\sum_{n=1}^{N_l}\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda_n^{'(1)})}\right)^2.
\end{aligned}\tag{C23}$$

Here we use a relation for $i=j$

$$\frac{1}{h_n}f_n^{i,i}(t)=\frac{1}{h_{n,t}}\delta_{t\in\mathcal{S}_n},\tag{C24}$$

where $h_{n,t}$ is a width of t in $\mathcal{S}_n$.

(ii) Plane but rapidly oscillating eigenvector $\psi_n^{(i)}$ with $C\simeq 0$, which is typical for the excited mode with $i>1$. In this case as was mentioned in the above a single summation $\sum_I\psi_n^{(i)}(I)$ is suppressed and (C16) has non-zero value only when $I=K$,

$$\begin{aligned}
&\frac{1}{h_n}\sum_I\sum_K \psi_n^{(i)\dagger}(I)\left\langle I\left|D_{N_5}^{-1}\right|J\right\rangle\left\langle J\left|\left(D_{N_5}^{-1}\right)^\dagger\right|K\right\rangle\psi_n^{(j)}(K)\\
&=\frac{1}{h_n}\sum_I \psi_n^{(i)\dagger}(I)\left\langle I\left|D_{N_5}^{-1}\right|J\right\rangle\left\langle J\left|\left(D_{N_5}^{-1}\right)^\dagger\right|I\right\rangle\psi_n^{(j)}(I)\\
&=\delta_{i,j}\left\langle I\left|D_{N_5}^{-1}\right|J\right\rangle\left\langle J\left|\left(D_{N_5}^{-1}\right)^\dagger\right|I\right\rangle,
\end{aligned}\tag{C25}$$

where I,J unsummed. X_l then becomes

$$\begin{aligned}
X_l(t)&=-\frac{a_5}{a^{10}}\sum_{n=1}^{N_l}\sum_{i>1}^{h_n}\frac{1}{h_n}f_n^{i,i}(t)\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda_n^{'(i)})}\right)^2\sum_{b,\beta}\left\langle I\left|D_{N_5}^{-1}\right|J\right\rangle\left\langle J\left|\left(D_{N_5}^{-1}\right)^\dagger\right|I\right\rangle\\
&=-\frac{a_5}{a^{10}}\frac{1}{4N_cn_{xyz}}\sum_{n=1}^{N_l}\sum_{i>1}^{h_n}\frac{1}{h_{n,t}}\delta_{t\in\mathcal{S}_n}\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda_n^{'(i)})}\right)^2
\end{aligned}$$

$$
\begin{aligned}
&\times \sum_{\vec{x}} \sum_{\alpha,a} \sum_{b,\beta} \left\langle \vec{x}, t, \alpha, a \left| D_{N_5}^{-1} \right| y, \beta, b \right\rangle \left\langle y, \beta, b \left| \left(D_{N_5}^{-1} \right)^{\dagger} \right| \vec{x}, t, \alpha, a \right\rangle , \\
&= \frac{1}{a_5} Y(t) \frac{1}{4 N_c N_x} \sum_{n=1}^{N_l} \sum_{i>1}^{h_n} \left(\frac{1}{2 \cosh \frac{N_5}{2} a_5 \widetilde{\lambda}(\lambda_n^{\prime (i)})} \right)^2 . \qquad \text{(C26)}
\end{aligned}
$$

In this case m_{5q} is given by the same formula as in the previous section with the completely localized eigenvector. In addition we assume that the spin and color eigenfunction is always rapidly oscillating and set their enhance factor to be unity as is discussed in the above.

The pion propagator in the denominator is expanded as

$$
\begin{aligned}
Y(t) &= \sum_{\vec{x}} \langle P(x) P(y) \rangle \\
&= -\frac{a_5^2}{a^{10}} \sum_{\vec{x},a,\alpha} \sum_{b,\beta} \Bigg(\sum_{n,m=1}^{N_l} \sum_{i,j=1}^{h_n} \frac{1}{h_n} \psi_n^{(i)}(I) \psi_m^{(j)}(I) \left\langle n, i \left| D_{N_5}^{-1} \right| J \right\rangle \left\langle J \left| (D_{N_5}^{-1})^{\dagger} \right| m, j \right\rangle \\
&+ \sum_{p,s,c} \sum_{k,s',d} \frac{1}{N_p} U_{\alpha}(p,s) U_{\alpha}^{\dagger}(k,s') e^{i(p-k)x} \left\langle p, s, a \left| D_{N_5}^{-1} \right| J \right\rangle \left\langle J \left| (D_{N_5}^{-1})^{\dagger} \right| k, s', b \right\rangle \Bigg) . \qquad \text{(C27)}
\end{aligned}
$$

By inserting a complete set $1 = \sum |I\rangle \langle I|$ the contribution from the localized eigenvectors is rewritten in term of $Y(t)$.

$$
\begin{aligned}
Y_l(t) &= -\frac{a_5^2}{a^{10}} \sum_{\vec{x},a,\alpha} \sum_{n,m=1}^{n_l} \sum_{i,j=1}^{h_n} \frac{1}{h_n} \psi_n^{(i)}(I) \psi_m^{(j)}(I) \sum_{b,\beta} \sum_{z,\gamma,c} \langle n, i | z, \gamma, c \rangle \left\langle z, \gamma, c \left| D_{N_5}^{-1} \right| J \right\rangle \\
&\times \sum_{w,\delta,d} \left\langle J \left| (D_{N_5}^{-1})^{\dagger} \right| w, \delta, d \right\rangle \langle w, \delta, d | m, j \rangle \\
&= -\frac{a_5^2}{a^{10}} \sum_{\vec{x},a,\alpha} \sum_{n,m=1}^{n_l} \sum_{i,j=1}^{h_n} \frac{1}{h_n} \psi_n^{(i)}(I) \psi_m^{(j)}(I) \\
&\times \sum_{b,\beta} \sum_{z,\gamma,c} \frac{1}{h_n} \psi_n^{(i)\dagger}(z,\gamma,c) \left\langle z, \gamma, c \left| D_{N_5}^{-1} \right| J \right\rangle \sum_{w,\delta,d} \left\langle J \left| (D_{N_5}^{-1})^{\dagger} \right| w, \delta, d \right\rangle \psi_m^{(j)}(w,\delta,d) \\
&= -\frac{a_5^2}{a^{10}} \sum_{\vec{x},a,\alpha} \sum_{b,\beta} \left\langle I \left| D_{N_5}^{-1} \right| J \right\rangle \left\langle J \left| (D_{N_5}^{-1})^{\dagger} \right| I \right\rangle_{I \in \cup_n \mathcal{S}_n} \\
&= \frac{\sum_{n=1}^{N_l} h_n}{4 N_c N_x} Y(t), \qquad \text{(C28)}
\end{aligned}
$$

where we take average for $I \in \cup_n \mathcal{S}_n$.

The contribution from the plane-wave modes is the same as that of the completely localized case by using the off diagonal property $\left\langle p, s, a \left| D_{N_5}^{-1} \right| n, i \right\rangle = 0$. Now we have a relation to write down $Y(t)$ in terms of $F(p)$ only,

$$
Y(t \to \infty) = -\frac{4 N_c N_x}{4 N_c N_x - h N_l} \frac{a_5^2}{a^{10}} n'_{xyz} \frac{4 N_c}{N_p} \sum_p F(p), \qquad \text{(C29)}
$$

where $h N_l \equiv \sum_{n=1}^{N_l} h_n$, and the anomalous quark mass m_{5q} can be written as

$$
\begin{aligned}
m_{5q} &= \frac{\lim_{t\to\infty} X(t)}{\lim_{t\to\infty} Y(t)} \\
&= \frac{1}{a_5}\frac{1}{4N_cN_x}\left(\sum_{n=1}^{N_l} 4^4 h_n \left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda_n^{\prime(1)})}\right)^2 + \sum_{n=1}^{N_l}\sum_{i>1}^{h_n}\left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}(\lambda_n^{\prime(i)})}\right)^2\right. \\
&\quad \left. +4N_c\sum_n \rho_n \left(\frac{1}{2\cosh\frac{N_5}{2}a_5\widetilde{\lambda}_n}\right)^2\right). \qquad \text{(C30)}
\end{aligned}
$$

In the actual case, the weight factor for the contribution from partially localized eigenstates lies between 1 and $4^4 h_n$ depending on the index i. Therefore we adopt $\tilde{h}_n^{(i)}$ such that $1 \le \tilde{h}_n^{(i)} \le 4^4 h_n$ as the weight factor in the text.

REFERENCES

[1] P. Ginsparg and K. Wilson, Phys. Rev. D25, 2649 (1982).

[2] M. Lüscher, Phys. Lett. B428, 342 (1998).

[3] P. Hasenfratz, Nucl. Phys. B525, 401 (1998).

[4] P. Hasenfratz, V. Laliena and F. Niedermayer, Phys. Lett. B427, 125 (1998).

[5] H. Neuberger, Phys. Lett. B417, 141 (1998); Phys. Lett. B427, 353 (1998); Phys. Rev. D57, 5417 (1998).

[6] Y. Kikukawa and T. Noguchi, hep-lat/9902022.

[7] R. Narayanan and H. Neuberger, Nucl. Phys. B412, 574 (1994); Nucl. Phys. B443, 305 (1995).

[8] D. Kaplan, Phys. Lett. B288, 342 (1992).

[9] Y. Shamir, Nucl. Phys. B406, 90 (1993).

[10] V. Furman and Y. Shamir, Nucl. Phys. B439, 54 (1995).

[11] T. Blum and A. Soni, Phys. Rev. D56, 174 (1997); Phys. Rev. Lett. 79, 3595 (1997); hep-lat/9712004.

[12] S. Aoki, T. Izubuchi, Y. Kuramashi and Y. Taniguchi, Phys. Rev. D62, 094502 (2000).

[13] For a review, see T. Blum, Nucl. Phys. B (Proc. Suppl.) 73, 167 (1999) and references there in.

[14] CP-PACS Collaboration, A. Ali Khan *et al.*, Phys. Rev. D63, 114504 (2001); Nucl. Phys. B (Proc. Suppl.) 83-84, 591 (2000).

[15] T. Blum, P. Chen, N. Christ, C. Cristian, C. Dawson, G. Fleming, A. Kaehler, X. Liao, G. Liu, C. Malureanu, R. Mawhinney, S. Ohta, G. Siegert, A. Soni, C. Sui, P. Vranas, M. Wingate, L. Wu and Y. Zhestkov, hep-lat/0007038; P. Chen, N. Christ, G. Fleming, A. Kaehler, C. Malureanu, R. Mawhinney, G. Siegert, C. Sui, P. M. Vranas, Y. Zhestkov, hep-lat/9812011; L. Wu *et al*, Nucl. Phys. B (Proc. Suppl.) 83-84, 224 (2000); G. T. Fleming *et al*, Nucl. Phys. B (Proc. Suppl.) 83-84, 363 (2000).

[16] H. Neuberger, Phys. Rev. Lett. 81, 4060 (1998).

[17] R. G. Edwards, U. M. Heller and R. Narayanan, Nucl. Phys. B540, 457 (1999); Phys. Rev. D59, 094510 (1999).

[18] T. DeGrand for MILC collaboration, Phys. Rev. D63, 034503 (2001).

[19] P. Hernandez, K. Jansen and M. Lüscher, Nucl. Phys. B552, 363 (1999).

[20] Y. Kikukawa, Nucl. Phys. B584, 511 (2000).

[21] I. Ichinose and K. Nagao, Nucl. Phys. B577, 279 (2000); hep-lat/0001030.

[22] R. C. Brower and B. Svetitsky, Phys. Rev. D61, 114511 (2000).

[23] M. Golterman and Y. Shamir, JHEP 0009, 006 (2000).

[24] F. Berruto, R. C. Brower and B. Svetitsky, hep-lat/0105016.

[25] S. Aoki, Phys. Rev. D30, 2653 (1984); Phys. Rev. Lett. 57, 3136 (1986); Nucl. Phys. B314, 79 (1989).

[26] S. Aoki and Y. Taniguchi, Phys. Rev. D59, 054510 (1999).

[27] S. Aoki, T. Kaneda and A. Ukawa, Phys. Rev. D56, 1808 (1997).

[28] R. G. Edwards, U. M. Heller and R. Narayanan, Nucl. Phys. B535, 403 (1998); Phys. Rev. D60, 034502 (1999).

[29] CP-PACS Collaboration, A. Ali Khan *et al.*, Nucl. Phys. B (Proc. Suppl.) 94, 725 (2001)

[30] A. Borici, hep-lat/9912040.

[31] CP-PACS Collaboration, in preparation.

[32] Y. Shamir, Phys. Rev. D62, 054513 (2000).
[33] H. Neuberger, private communication.
[34] M. Lüscher, private communication.
[35] S. Nakamura, private communication. W. Kirsch, “Random Schrödinger operators”, in Lecture Notes in Physics 345 (Springer-Verlag).
[36] P. W. Anderson, Rev. Mod. Phys. 50, 191 (1978).
[37] F. Berruto, R. Narayanan and H. Neuberger, Phys. Lett. B489, 243 (2000).
[38] R. G. Edwards and U. M. Heller, Phys. Rev. D63, 094505 (2001).
[39] P. Hernandez, K. Jansen and M. Lüscher, hep-lat/0007015.
[40] R. B. Laughlin, Phys. Rev. B23, 5632 (1981).

FIGURES


g^2
∞
$\langle\bar{\psi}i\gamma_5\tau^3\psi\rangle \neq 0$
$\langle\bar{\psi}i\gamma_5\tau^3\psi\rangle = 0$
A
B
A
$M_c(g^2)$
$M_c(g^2)$
g_c^2
A
A
A
A
0
M
0
2
4
6
8

FIG. 1. The expected phase structure of lattice QCD with the Wilson fermion in the M–g^2 plane. In the region A, $\langle\bar{\psi}i\gamma_5\tau^3\psi\rangle = 0$ (the parity-flavor symmetric phase), while $\langle\bar{\psi}i\gamma_5\tau^3\psi\rangle \neq 0$ (the parity-flavor broken phase) in the region B.



FIG. 2. $F_c(p)$ as a function of $\widetilde{\lambda}(p)$ at $M = 0.8$(left) and $M = 1.4$(right). Both F_c and $\widetilde{\lambda}$ are evaluated on $16^3 \times 32$ momentum lattice, where the periodic boundary condition is adopted for spatial direction and anti-periodic boundary condition is set for temporal direction in order to prevent a singularity at the origin.



FIG. 3. Three types of $\rho(\lambda)$ adopted in calculation of m_{5q}. The left one corresponds to type (1) with $\delta < 0$, the center is type (2) with $\delta = 0$ and the right is type (3) with $\delta > 0$.



FIG. 4. m_{5q} as function of N_5 evaluated with three types of $\rho(\lambda)$. The left one corresponds to type (1) with $\delta = -0.5$, the center is type (2) with $\delta = 0$ and the right is type (3) with $\delta = 0.5$.